

Vultures In Gidh Karai, Jhalawar, Rajasthan, India: Threats, Recommendations and Conservation Measures

Taruna Patidar¹, Dr. Surabhi Shrivastava²

¹Researcher Wildlife Science

²Former HOD, Coordinator, Department of Wildlife Science, University of Kota, Kota, Rajasthan, India

Abstract

Vultures used to dominate the environment as the most frequent scavengers prior to the 1980s, but their numbers catastrophically dropped to >95%. A veterinary NSAID named Diclofenac, which was nephrotoxic to birds and caused renal failure and avian gout, was employed in this situation. Diclofenac was outlawed in 2006 by the governments of India, Pakistan, and Nepal in an effort to preserve the declining vulture population. The development of non-toxic meloxicam as a substitute for diclofenac has led to the stabilisation of Gyps vulture populations in various regions of Southern Asia. There are nine different species of vultures in India (both resident and migratory), three of which are categorised as Critically Endangered on the IUCN Red List of Threatened Species and are thus given the strictest legal protection under Schedule-I of the Indian Wildlife Protection Act. However, Rajasthan is home to seven different vulture species. The study area, i.e., Gidh Karai, located in the Jhalawar district of Rajasthan (India), is an ideal location for breeding and roosting Indian and Egyptian vultures. And this study briefly highlights the underlying threats, recommendations and conservation measures that need to be implemented for the protection of these vultures. It is also a sincere effort to draw attention towards the crisis of vultures and the overall need for conservation.

Keywords: Conservation, Gidh Karai, recommendations, threats

1. Introduction

Vultures, now on the verge of extinction, ultimately pose a sincere concern for their conservation. According to Ogada et al. (2012), currently fourteen vulture species (61%) are threatened with extinction worldwide, and the most rapidly declining curves occur in the vulture-rich regions of Asian and African countries. Of the nine species of vultures recorded in the Indian subcontinent, five belong to the genus *Gyps* (Ali and Ripley, 1995; Prakash, 1999; Gadhvi and Dodia, 2006).

Of the total recorded species of Gyps vultures in India, three species, namely *Gyps bengalensis* (White-rumped Vulture), *Gyps indicus* (Indian Vulture/Long-billed Vulture) and *Gyps tenuirostris* (Slender-billed Vulture), are resident species, while the other two, i.e., *Gyps himalayensis* (Himalayan Vulture) and *Gyps fulvus* (Griffon Vulture), are migratory or winter visitors (Prakash et al., 2003). *Gyps indicus* (Sibley and Monroe, 1990; 1993) is divided into two species: *G. indicus* and *G. tenuirostris* (Rasmussen and Parry, 2001).

Vultures, being birds of prey, have highly specialised musculature in their feet, though this has now degenerated to some extent due to changing scavenging patterns or ways of life.

The Old World vultures had a comfortable population status until a couple of decades ago, but it changed radically and by the year 2015, eight out of 16 species were considered Critically Endangered (revealing the possibility of their extinction within 10 years), three were endangered, four were Near threatened and only one was of Least Concern. There was a likely and feasible chance that several of these species could become extinct in the near future unless effective conservation measures were implemented in their distribution range. The major reason for the rapid decline in the population of vultures is mainly due to the poisoning of their food, both intentional and unintentional. Due to sincere and optimistic conservation efforts in Europe, the populations of some species like the Egyptian Vulture, Bearded Vulture and Cinereous Vulture have shown observable recovery (Margalida and Oliva-Vidal, 2017). Concerns including habitat loss or degradation (Carette et al., 2007), food availability, collisions, and electrocution by electrical power lines (Anderson and Kruger, 1995; Boshoff et al., 2011) that work differently in different parts of the world are some other risks to vultures.

Rajasthan has one of the largest surviving populations of vultures in India. Of the nine species of vultures found in different habitats in India, seven occur in the Indian state of Rajasthan (Naoroji 2007; Ali & Ripley 2007). Thus, it makes the existential vulture breeding colonies worthy of virtuous protection.

1. Review of Literature

- Shultz *et. al* (2004) showed that the observed high proportions of dead *Gyps bengalensis* and *Gyps indicus* carcasses with visceral gout or diclofenac residues were broadly identified in India and Nepal, as indicated in the associated post-mortem findings.
- Prakash *et. al* (2007) surveyed on identified tracks in 2007 to repeat surveys done previously in 1992, 2000, 2002 and 2003 to determine the population trend in three species of vultures and also to get the surviving population trend of vultures in 2007. The results revealed that the vulture population of the three species decreases continually at an alarming rate. The numbers of Oriental white-backed vultures decreased by 99.9% between 1992 and 2007. The analogous decline in the integrated total of *Gyps indicus* and *G. tenuirostris* was 96.8%. The Oriental white-backed vultures' average population declined by 43.9% annually between 2000-2007, whereas the *G.indicus* and *G. tenuirostris* combined average populations declined by over 16% annually. A complete ban on the use of diclofenac in livestock treatment and the establishment of conservation breeding centres were suggested to prevent the extinction of these three species of vultures.
- Chhangani (2007) studied 5,000 vultures of seven species in the Rajasthan and Gujarat states of India. He concluded that besides the NSAID diclofenac, other reasonable causes are leading to vulture decline, such as habitat loss due to huge mining, quarrying, blasting and logging operations in Rajasthan and Gujarat. Other reasons include the unavailability of food and water, predation or attacks by wild dogs and collisions or accidents. Changes in agriculture and livestock rearing land-use practises have also affected the vulture's breeding ecology. He also imbibes the importance of studying the genetic diversity of all *Gyps* vultures in India and conducting extensive surveys to ascertain the decline and other issues related to their biology.
- Jha (2015) studied three seasons to evaluate the abundance and diversity of vultures in Uttar Pradesh during 2010–11. Of the nine species found in India, six reside in UP. The counts and qualitative

evaluations showed that the vulture population had decreased during the previous 10 to 15 years, with the primary causes being the use of diclofenac, a lack of food, and habitat destruction.

- Dead carcasses were eliminated on the village outskirts, where crows, egrets and dogs competed with vultures. He suggested effective conservation by securing safe and sufficient food, recovery from accidents and rehabilitation, and a protected environment for the vultures in UP.
- Rao (April 2020) found that the vulture population rose by 103% in Bundelkhand in the past decade. However, he also discussed the factors that led to the population decrease, mainly the destruction of forests and disruption of nesting sites. The stray cattle also raise a subsequent concern of invincible effects on food and habitat for vultures. He also highlights the threats to vulture species and the reviving of existential vulture restaurants.

2. Study Site




Fig. 1: Location of the Vultures on the Bank of the Kali-Sindh River

The chosen site is situated in the Jhalawar Forest Division's Khanpur Range, nearby Hichar Village. Mukundara Hills Tiger Reserve (MHTR) is located in close proximity to Gidh Karai. The vulture colonies are situated at Latitude 24.656049 and Longitude 76.1821 on the Gidh Karai cliffs beside the Kali-Sindh River. On the Kali-Sindh river bank, vulture colonies can be seen on each side of the cliffs.

3. Materials & Methods

Field Survey:

The field survey was conducted at a location called Gidh Karai, where colonies of Indian and Egyptian vultures were found. It comprised a 3-kilometre hike to the location on the banks of the Kali Sindh River from the Gagon fort (a hill water fort located 10km approx. in the north-eastern part of Jhalawar) and Belinda Ghat. From the other side, it is adjacent to Harish Chandra Sagar Pariyojana at Hichar Village (Jhalawar). Vultures have the natural protection they need to carry out their roosting, nesting and breeding activities in the colonies, which are present on the cliffs on both sides of the Kali Sindh River.

4. Observations & Results

Threats to Vultures

India is home to the world's greatest number of cattle. And since vultures are scavengers, they like to eat dead and decaying bodies. Their long, bare necks and lack of feathers allow them to plunge deep into the carrion in search of the inner, softer components. Their efficiency is well-marked, as they can finish a deer or a herd of cattle in less than 20 minutes. However, their natural scavenging behaviour has now become a concern for their survival. To prevent the spread of disease, it is now preferred to adapt the carcass disposal technique by dumping them later. As a result, the vulture population is threatened by a subsequent decline in food availability.

The effect of administering diclofenac, a veterinary non-steroidal anti-inflammatory drug, to cattle to treat fever, inflammation, and pain-related illnesses or infections is harmful. It was found to be fatal even in minor dosages, causing renal failure and uric acid accumulation in the blood, as well as crystallisation around internal organs, a disease known as visceral gout.

Vultures are the most vulnerable to human meddling or activity. Illegal hunting, power line collisions and electrocution, cliff mining or quarrying, forest degradation through logging or tree felling, livestock grazing, and disturbance by langurs owing to interruptions in their regular routines are all factors that are displacing vultures. In areas of their natural environment, they are threatened by habitat loss. They are slow breeders, laying a single egg during one breeding season, and if their nests are disturbed, they are unlikely to build nests in the same location.

When a farmer or livestock owner poisons the bodies of deceased domestic animals to eliminate a rogue predator, the vultures become victims of retaliatory or inadvertent poisoning. This event occurs occasionally and is not responsible for the population crash. However, the vulture population is so low, that even such operations could result in local extinctions and should be avoided.

Insecticides and pesticides that damage the environment and accumulate in water bodies serve as a possible contamination source, resulting in significant losses of these birds due to bio-magnification.

Infectious pathogens are another danger to vultures. A post-mortem examination of 28 Indian vulture and White-rumped vulture carcasses collected across India reveals signs of infectious disease (Cunningham *et al.*, 2003).

5. Summary & Conclusion

Why is conserving the present study area important?

We may conclude from field observations that there is a significant amount of the immature population in this location. This emphasizes the necessity of preserving the area so that critically endangered Indian vultures and endangered Egyptian vultures can continue to thrive in their native habitat.

Nesting takes place during the winter months, from November to February. Therefore, limiting people's movement and eliminating external disruptions in this location will help the vultures overcome their psychological dread.

There is a good record of Indian Gyps vultures and Egyptian vultures at the site. Crocodiles in the Kali Sindh River and Hanuman langurs are also seen.

Conserving them offers significant benefits for the Jhalawar region's environment, including:

Carrion acts as a breeding environment for potentially harmful bacteria, perhaps leading to direct or indirect infections, and is a source of diseases such as anthrax. Therefore, an increase in uneaten carcasses in adjacent settlements offers a direct hazard to human health. Vultures clean the environment

and safeguard humans, livestock, and wildlife from diseases and sickness by quickly and efficiently eliminating carcasses.

Wider environmental consequences could include pollution from rotting carcasses and the spread of illnesses such as rabies as the feral dog population expands.

6. Recommendations

Measures for the conservation of vultures

Based on the field observations, the following measures can be taken:

1. Eliminating the principal causal agent, diclofenac, from the environment by preventing diclofenac from entering the veterinary sector through human formulations.
2. Designating the region as a Conservation reserve will provide legal protection for the vulture site while also restricting human mobility in the area.
3. Submitting the location to the State Forest Department for inclusion in Vulture Safe Zones (VSZ), which have been recommended to be established in various sections of the country to protect the country's remaining vulture population.
4. Collecting dead vultures to determine the reasons for their deaths in order to monitor vulture conservation and recovery.
5. Establishing and expanding a vulture breeding and care centre
6. Preventing more deaths by keeping a careful eye on unlawful mining in these areas.
7. Raising awareness, particularly among veterinary formulation users, to decrease the impact on vultures.
8. Monitoring the implementation of the "Vulture Conservation Action Plan 2020-2025."
9. DECLARATION OF THIS AREA AS A VULTURE SAFE ZONE:

After two years of searching, a VSZ could be established, ensuring that:

1. No Diclofenac or visceral gout was identified in deceased vultures in the area, and
2. Vulture numbers within the VSZ are either steady or expanding.
3. Vultures can also be found in Mukundara Hill Tiger Reserve's surrounding locations. In the future, appropriate research can be conducted in that location, and the site can be preserved for the Vultures.

7. Future Action Plan at the State Level

• The Role of State Forest Departments

All PCCFs (Wildlife) and CWLWs will be responsible for executing the APVC 2020–2025, according to the Action Plan for Vulture Conservation (APVC).

• Setting Up State Committees

The MoEFCC will instruct state forest ministries to form state vulture conservation committees to carry out the NC's recommendations and instructions. The State Committee will be composed of the Forest Secretaries of all state governments, with the Chairman being the PCCF and CWLW, and the members being the Director General/Director, Animal Husbandry, and Drug Controller. The BNHS will be a non-official member, and the Member Secretary will be CCF (Wildlife) HQ.

The committee will oversee the state's implementation of APVC 2020–2025. The responsible agency will be in charge of carrying out the NC's decision.

8. Vulture Conservation in India

Vulture Conservation Breeding Programme

The programme began in 2004 to conserve the three resident Gyps vulture species, the White-backed vulture, Long-billed Vulture, and Slender-billed Vulture, from extinction as a result of the 2006 Action Plan for Vulture Conservation (MoEF, Government of India 2006).

Its goal is to raise birds in captivity under near-natural conditions.

The first Vulture Conservation Breeding Centre was constructed in Pinjore, Haryana, in 2004 due to the combined efforts of the Haryana Forest Department and the BNHS. It also ensures the health of the birds before they are released back into the wild.

Vulture Conservation Breeding Centres (VCBC) have been established in eight locations across the country: Rajabhatkhawa, Buxa National Park (West Bengal) in 2006; Rani (Assam) in 2008. The Central Zoo Authority (CZA) collaborated with state zoos to establish five new centres in Van Vihar, Bhopal (Madhya Pradesh); Nandankanan Zoo, Bhubaneswar (Odisha); Nehru Zoological Park, Hyderabad (Telangana); Muta Zoo, Ranchi (Jharkhand); and Sakkarbaug Zoo, Junagadh (Gujarat).

Now, the most recent Action Plan for Vulture Conservation 2020–2025 (APVC) is working to conserve our vultures on the ground.

Action Plan for Vulture Conservation 2020-2025

Vital aspects:

Drug control:

- a) To prohibit the use of veterinary NSAID Diclofenac.
- b) Restrict the vultures' food poisoning, the cattle carcasses, with veterinary NSAIDs through safe disposal of carcasses and the regulation of proper sale, distribution and administration of drugs. It also assures that the livestock is medicated by qualified veterinarians.
- c) To facilitate vulture safety testing and trials of available veterinary NSAID compounds. And to ensure that only the safest ones are let onto the market.
- d) To ensure this, the DGCI must design a system that will automatically remove a veterinary medicine from use if it is discovered to be dangerous to vultures. It would ensure that harmful medications like aceclofenac and ketoprofen, as well as diclofenac, are no longer available on the market.

9. Conservation Measures

- a) Enhancement of the country's conservation breeding programme.
- b) There are now only eight Vulture Conservation Breeding Centres in the country, and there is a growing demand for more Vulture Conservation Breeding Centres in various parts of the country. These centres' primary goal is the breeding of vultures, but they can also serve as Vulture Conservation Centres, where samples and data acquired in the wild can be analysed and stored. The Breeding Programme also includes the Red-Headed Vulture and Egyptian Vulture species.
- c) To enforce the Vulture Safe Zone Programme in eight distinct locations across the country with established vulture populations. To campaign for and proclaim at least one Vulture Safe Zone in each state, as well as to continue working on the existing efforts of Vulture Safe Zone.
- d) Monitoring of vultures periodically across the country.

- e) Other causes of death, apart from veterinary medicine poisoning of vulture food, should be considered.
- f) Four rescue centres have been created for distinct geographical locations, including Pinjore in the north, Guwahati in the northeast, Bhopal in central India, and Hyderabad in the south.

Other Efforts:

o National:

- The Vulture Conservation Breeding Programme was established by CZA and BNHS in collaboration. For the breeding and conservation of Indian vulture species, a Jatayu Conservation Breeding Centre (Asia's first vulture breeding facility) has been established at Pinjore. In 2001, the Vulture Care Centre (VCC) was founded to investigate the causes of vulture fatalities.
- MoEFCC is in charge of implementing the Action Plan for Vulture Conservation 2020–2025 (APVC), which includes the combined efforts and active cooperation of MoH&FW and MoAHDF for vulture conservation in India. To some extent, the Ministries of Power, Renewable Energy, Chemicals, and Fertilizer will also need to be incorporated. As a result, it would necessitate cross-ministerial cooperation, making the conservation of vultures far more difficult than that of any other wild species.

o International: SAVE (Saving Asia's Vultures from Extinction)

The consortium of 24 organisations that establish virtuosity in vulture conservation was established in 2011 that coordinates conservation, fund-raising and campaigning to help save vultures. Its objective is to promote worthy action to prevent the extinction of Gyps vultures in South Asia.

International Action Plan

The APVC benefits from international programmes such as the Vulture MsAP (Multi-Species Action Plan to Conserve African-Eurasian Vultures), produced by the Conservation of Migratory Species (CMS) Raptors MoU, SAVE & National Action Plan, and the Vulture Conservation Action Plan for Nepal 2015–19. CMS Parties adopted it at COP12, which took place in Manila, Philippines, from October 23 to October 28, 2017. All nine species of vultures found in India are included in the action plan.

Acknowledgement

I would like to express my sincere thanks to the Jhalawar Forest Department, especially Mr. Arunkumar Devaraju (IFS), my supervisor (Dr. Surabhi Shrivastava) and my family for their constant support and motivation to complete the research work. I would like to extend my special regards to Dr. Vikram Thak and Mr. Deepak Jadav, who helped me collect the information.

References

1. Yadav, V. K. & Chauhan, P. S. (2018). Avifaunal Diversity and Status of Jhalawar Forest Division, South-Eastern Rajasthan, India. *Indian Journal of Ecology*, 45(1): 107-116.
2. Swan, G.; Naidoo, V.; Cuthbert, R.; Green, R.E.; Pain, D.J.; Swarup, D.; Prakash, V.; Taggart, M.; Bekker, L.; Das, D. and Diekmann, J. (2006). Removing the threat of diclofenac to critically endangered Asian vultures. *PLoS biology*, 4(3);e66.

3. Suthar, S. *Assessment and Management Strategies of Vultures in Hadoti Region of Rajasthan* (Doctoral dissertation, College of Horticulture and Forestry, Jhalawar).
4. Suthar, S.; Sharma, A.; Chauhan, P. S.; Chauhan, K.; Nagar, B. & Maurya, I. B. (2020). Assessment of Diversity, Population and Related Threats to Vultures in Hadoti Region of Rajasthan. *Int. J. Curr. Microbiol. App. Sci*, 9(12): 1302-1309.
5. Shultz, S.; Baral, H.S.; Charman, S.; Cunningham, A.A.; Das, D.; Ghalsasi, G.R.; Goudar, M.S.; Green, R.E.; Jones, A.; Nighot, P. and Pain, D.J. (2004). Diclofenac poisoning is widespread in declining vulture populations across the Indian subcontinent. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271(suppl_6):S458-S460.
6. Patidar, T. and Shrivastava, S. (2023). Study of Population Status of Vultures in Gidh Karai, Jhalawar, Rajasthan. *International Journal of Research in Academic World*, 2(5):88-93
7. Patidar, T. and Shrivastava, S. (2023). Study of habitat suitability factors favouring the survival of vultures in Gidh Karai, Jhalawar, Rajasthan, India. *International Journal of Science and Research*, 12(5):2426-2431
8. Reinde, M. (September 3, 2019). Poison Pill: The Mysterious Die-Off of India's Vultures. *DISTILLATIONS, Science History Institute*. Available at: <https://www.sciencehistory.org/distillations/poison-pill-the-mysterious-die-off-of-indias-vultures>.
9. Rasmussen, P.C. and Parry, S.J. (2001). The taxonomic status of the long-billed vulture *Gyps indicus*. *Vulture News*, 44:18-21.
10. Rao, M. M. (2020). Vultures in Bundelkhand rise, but stray cattle and other concerns persist. *MONGABAY*. <https://india.mongabay.com/2020/04/vultures-in-bundelkhand-rise-but-stray-cattle-and-other-concerns-persist/>
11. Prakash, V. (1999). Status of vultures in Keoladeo National Park, Bharatpur, Rajasthan, with special reference to population crash in Gyps species. *Journal Bombay Natural History Society*, 96:365-378.
12. Prakash, V.; Pain, D.J.; Cunningham, A.A.; Donald, P.F.; Prakash, N.; Verma, A.; Gargi, R.; Sivakumar, S. and Rahmani, A.R. (2003). Catastrophic collapse of Indian white-backed Gyps bengalensis and long-billed Gyps indicus vulture populations. *Biological conservation*, 109(3):381-390.
13. Prakash, V.; Green, R.E.; Pain, D.J.; Ranade, S.P.; Saravanan, S.; Prakash, N.; Venkitachalam, R.; Cuthbert, R.; Rahmani, A.R. and Cunningham, A.A. (2007). Recent changes in populations of resident Gyps vultures in India. *Journal of the Bombay Natural History Society*, 104(2):129-135.
14. Prakash, V.; Bishwakarma, M.C.; Chaudhary, A.; Cuthbert, R.; Dave, R.; Kulkarni, M.; Kumar, S.; Paudel, K.; Ranade, S.; Shringarpure, R. and Green, R.E. (2012). The population decline of Gyps vultures in India and Nepal has slowed since veterinary use of diclofenac was banned. *PLoS One*, 7(11):e49118.
15. Official website of Jhalawar district - <https://www.jhalawar.rajasthan.gov.in/>
16. Oaks, J.L.; Gilbert, M.; Virani, M.Z.; Watson, R.T.; Meteyer, C.U.; Rideout, B.A.; Shivaprasad, H.L.; Ahmed, S.; Chaudhry, M.J.I.; Arshad, M. and Mahmood, S. (2004). Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature*, 427(6975):630-633.
17. Naoroji, R. & Schmitt, N. J. (2007). *Birds of prey of the Indian subcontinent*. Om Books International.
18. Gadhvi, I. R. & Dodia, P. P. (2006). Indian White-backed Vultures *Gyps bengalensis* nesting in Mahuva, Bhavnagar district, Gujarat, India. *Indian Birds*, 2(2): 36.

19. Devaprakash, J. (2016). Fall of the Vultures.
20. Chhangani, A. K. (2007). Why are Vultures Dying? *DownToEarth*. <https://www.downtoearth.org.in/blog/why-are-vultures-dying-7054>
21. Chhangani A.K. and Mohnot S.M. (2004). Is diclofenac the only cause of vulture declines? *Current Science*, 87(11): 1496-1497.
22. Chaudhry, M. J. I.; Ogada, D. L.; Malik, R. N.; Virani, M. Z. & Giovanni, M. D. (2012). First evidence that populations of the critically endangered Long-billed Vulture *Gyps indicus* in Pakistan have increased following the ban of the toxic veterinary drug diclofenac in south Asia. *Bird Conservation International*, 22(4): 389-397.
23. Carrete, M.; Grande, J. M.; Tella, J. L.; Sánchez-Zapata, J. A.; Donázar, J. A.; Díaz-Delgado, R. & Romo, A. (2007). Habitat, human pressure, and social behavior: Partialling out factors affecting large-scale territory extinction in an endangered vulture. *Biological Conservation*, 136(1): 143-154.
24. Boshoff, A.F.; Minnie, J.C.; Tambling, C.J. and Michael, M.D. (2011). The impact of power line-related mortality on the Cape Vulture *Gyps coprotheres* in a part of its range, with an emphasis on electrocution. *Bird Conservation International* 21: 311-327. doi: 10.1017/S095927091100013X
25. Anderson, M. D. & Kruger, R. (1995). Power line electrocution of eighteen African White-backed Vultures. *Vulture News*, 32(1): 18.
26. Ali, S. and Ripley, S.D. (1995). *A Pictorial Guide to the Birds of Indian Subcontinent*. Bombay.
27. Ali, S. and Ripley, S.D. (2007). *Handbook of the Birds of India and Pakistan*. Bombay Natural History Society and Oxford University Press. Compact ed.
28. Ogada, D. L.; Keesing, F. & Virani, M. Z. (2012). Dropping dead: causes and consequences of vulture population declines worldwide. *Annals of the New York Academy of Sciences*, 1249(1): 57-71.
29. Ministry of Environment, Forest and Climate Change, MoEF&CC (2020). *Action Plan for Vulture Conservation in India, 2020-2025*. <https://save-vultures.org/wpcontent/uploads/2020/11/20-11-India-National-Vulture-Action-Plan-2020-25.pdf>.
30. Monroe Jr., B.L. & Sibley, C.G. (1993). *A World Checklist of Birds*. Yale University Press, New Haven and London: [I]-XIX, 1-393.
31. Margalida, A. & Oliva-Vidal, P. (2017). The shadow of diclofenac hangs over European vultures. *Nature ecology & evolution*, 1(8): 1050-1050.
32. Jha, K.K. (2015). Distribution of vultures in Uttar Pradesh, India. *Journal of Threatened Taxa*, 7(1):6750-6763.
33. Google Earth Pro (2021). Rajasthan, India.
34. Green, R. E.; Newton, I. A. N.; Shultz, S.; Cunningham, A. A.; Gilbert, M.; Pain, D. J. & Prakash, V. (2004). Diclofenac poisoning as a cause of vulture population declines across the Indian subcontinent. *Journal of Applied ecology*, 41(5): 793-800.