

# Neuromuscular Diseases in Intensive Care

Yahya Lahrichi<sup>1</sup>, Ali Derkaoui<sup>2</sup>, Ibrahim Bechri<sup>3</sup>, Abdelkrim A Shimi<sup>4</sup>,  
Mohammed Khatouf<sup>5</sup>

<sup>1</sup>Intern, Anesthesiology and Intensive Care Department A1, Hassan II University Hospital, Fez, Morocco, Faculty of Medicine and Pharmacy of Fez, Sidi Mohammed Benabdellah University, Fez, Morocco

<sup>2,3,4,5</sup>Professor, Anesthesiology and Intensive Care Department A1, Hassan II University Hospital, Fez, Morocco, Faculty of Medicine and Pharmacy of Fez, Sidi Mohammed Benabdellah University, Fez, Morocco

## Abstract:

Guillain-Barre syndrome and myasthenia gravis are common in the intensive care unit (ICU), and are not always easy to diagnose in the emergency department. The short-term risk is the occurrence of respiratory muscle paralysis, which must be monitored by repeated measurement of vital capacity, and the existence of respiratory impairment requires transfer to intensive care. Apart from specific therapeutic options (plasma exchange, intravenous immunoglobulins and anticholinesterase agents), management is mainly symptomatic. Our work is a retrospective study, concerning 23 cases of neuromuscular diseases in the ICU from a period from 2018 to 2022. The various epidemiological, Clinical, paraclinical, Therapeutic and evolutionary data were collected from the files of the intensive Care Department A1 of the Hassan II University Hospital of Fez.

**KEYWORDS:** respiratory distress- immunoglobulins- nerve stimulation- myasthenia gravis-Guillain-Barré syndrome- nosocomial pneumonia

## Introduction:

- Guillain-Barré syndrome (GBS) has become, since the eradication of acute anterior poliomyelitis, the main cause of extensive acute paralysis in developed countries. Given the risk of short-term severity, it is a neurological emergency. Diagnosis is generally clinical, with additional tests used mainly to rule out a differential diagnosis. Symptomatic treatment is paramount, aimed at preventing complications of motor deficit. (1,2).

- Myasthenia is a chronic autoimmune disease linked to a defect in the transmission of nerve impulses between the motor nerve and striated muscle. A disease of the neuromuscular junction due to blockage of motor plate receptors by anti-acetylcholine receptor antibodies. It manifests as muscular weakness that worsens with exertion and improves with rest. Diagnosis is straightforward, but treatment is more difficult due to the unpredictable course of the disease. Its severity lies in the risk of respiratory complications that can be life-threatening. (3,4).

**Materials and methods:**

The aim of our study is to describe the clinical, diagnostic, therapeutic and evolutionary aspects of neuromuscular diseases in the intensive Care Department A1 of the Hassan II University Hospital of Fez between 2018 and 2022. Our work is a retrospective study, containing 23 cases from a period from 2018 to 2022. The various epidemiological, clinical, paraclinical, therapeutic and evolutionary data were collected from the files of the intensive Care Department A1 of the Hassan II University Hospital of Fez.

**Results:**

-During our study 23 cases were included. (Table 1) The mean age was 49.72, with extremes ranging from 17 to 81 years (Figure 1), and males predominated. An infectious syndrome was present in 45% of cases (Figure 2). The main symptom (80% of cases) was heaviness of the limbs. Respiratory distress was reported in 8 patients, swallowing disorders in 6 and sphincter disorders in 3. Only one patient was admitted in a state of cardiocirculatory shock of hypoxic origin (Sd de guillanbarré diagnosed by EMG) and another for a myasthenic crisis. The electroneuromyogram showed 3 types of electromyography (demyelinating in 52% of cases, axonal and post-synaptic neuromuscular block in 4%). In addition to symptomatic treatment, which was the norm in all our patients depending on the type of detesis, specific treatment was instituted depending on the type of pathology: plasmapheresis in 17.39% of cases, immunoglobulins in 79%, anticholinesterase agents and corticosteroids in 13.04% of cases (Table 2). The evolution was marked by complications especially of infectious order made nosocomial pneumonia in 50% of the cases ,the mortality in our series was 21,73% of the cases .The majority of patients died in respiratory distress.

**Discussion:**

Guillain-barré syndrome and myasthenia gravis are neuromuscular diseases of variable incidence, estimated at between 0.81 and 1.89 per 100,000 inhabitants per year in Western countries(1,2). Myasthenia classically affects women between the ages of 20 and 30, which is consistent with the results of our series(3,4) . Diagnosis is predominantly clinical, with additional tests performed mainly to rule out a differential diagnosis. GBS evolves in three phases: an extension phase, a plateau phase and a recovery phase; generally progressive weakness of more than one limb associated with areflexia evolving in less than 4 weeks (5).Heaviness of the limbs was found in 18 patients with major respiratory distress in 30% of cases. The technique most commonly used to assess neuromuscular transmission is repeated nerve stimulation. Stimulation at 2 to 5 Hz results in a progressive decrement of the muscle component of the action potential in the myasthenic patient (from at least 10% between 1st and 5th stimulation). This test is abnormal in 75% of patients with generalized myasthenia but in less than 50% of patients with a purely ocular form, and post-synaptic neuromuscular block was found in only one patient in our series. Polyvalent immunoglobulins are as effective as plasma exchange in terms of mortality and clinical recovery (weaning from mechanical ventilation, recovery of walking) in the short (four weeks) and long (one year) term. (6,7,8).Immunoglobulins were administered to 18 patients (78.26%).We had 4 deaths out of a total of 20 patients hospitalized for Guillain-Barré syndrome, i.e. 25% of cases, a high percentage given the limited number of cases in our series. The reported mortality for Guillain-Barré syndrome is 6% in the article by Doets et al. 2018 (9).Tolerance is reported to be better, with fewer discontinuations than with plasma exchange. IV immunoglobulins were administered in 18 of our patients. Risk factors for mortality were: age, nadir disease severity, need for mechanical ventilation and hyponatremia.

**Conclusion:**

Patients with neuromuscular diseases are at greater risk of ICU admission, especially those with chronic respiratory failure, cardiac impairment, reduced mobility or swallowing disorders. Early detection and prompt initiation of appropriate treatment can improve morbidity and mortality in these patients.

**Reference**

1. Gwénolé Abgrall. Epidemiology of intensive care neuromyopathy in Fort-de-France: incidence, facilitating factors and external validation of a predictive score: the NERVIF study . Human Medicine and Pathology. 2015.
2. Jean-Romain Garric. Guillain-Barré and Strohl syndrome in intensive care. Sciences du Vivant [q-bio]. 2014
3. Svahn J, Chenevier F, Bouhour F, Vial C. Myasthenias and myasthenic syndromes. EMC - Neurologie 2018;
4. BENCHEKROUN EM. La myasthénie: Expérience du service de neurologie de l'hôpital militaire Moulay Ismaël de Meknès. ThèseMed : Université Sidi Mohammed Ben Abdellah: 2016
5. T. Ritzenthaler, T. Sharshar, D. Orlikowski. Guillain-Barré syndrome. EMC - Anesthesia-Reanimation volume 11, n°3, July 2014.
6. Dr Khaoula BALILI Management of Guillain-Barré syndrome: comparison between Intravenous Immunoglobulins and Plasmapheresis These FMPM 2020.
7. Grapperon A, Attarian S. Guillain-Barré syndrome. EMC - Neurol. 2016;
8. Bouche P, Léger JM, Vallat JM. Peripheral neuropathies. Polyneuropathies and multiple mononeuropathies. Volume 2. Rueil-Malmaison:Doin; 2004.
9. Doets, A. Y., Verboon, C., van den Berg, B., Harbo, T., Cornblath, D. R., Willison, H. J., ... Bateman, K. Regional variation of Guillain-Barré syndrome. Brain.(2018).

Number of patients hospitalized	2819
Number of patients with Guillain-Barré syndrome	20
Number of patients with myasthenia gravis	3

**Table 1: Incidence of neuromuscular diseases in our series**




Figure 1: Age distribution




Figure 2: Type of trigger

Type	Workforce	Percentage
Plasmapheresis	4	17.39%
Immunoglobuline	18	78.26%
Anticholinesterase	3	13.04%
Corticosteroidtherapy	3	13.04%

Table 2: Specific treatment in our series