

An Observational Study to Find Impact of Covid19 Infection on Causation and Prognosis of Pancreatitis

Neeraja Kattula

Asst Prof of Zoology

Department of Zoology

BRR Govt Degree College, Jadcharla, Telangana, India-509301

Abstract

Objective: To find impact of covid19 infection on causation and prognosis of pancreatitis

Design: Prospective observational study

Setting: A large, academic, tertiary medical center

Methodology : Clinical and laboratory data of all consecutive patients with a primary diagnosis of TS during past 6months with history of covid 19 infection was collected .Clinical and radiological grading was taken into consideration for finding severity.

Results: out of 30 patient 4 patients had severe covid 19 infection requiring more than 10days of hospitalization but though 1 of those 4 patient developed recurrent pancreatitis but severity score was less(clinical and radiological).14 had mild covid 19 infection with minimal respiratory and more of GI symptoms .Out of these 14 ,10 developed severe acute necrotizing pancreatitis requiring ICU stay and prolonged hospital stay .12 patients had mild covid 19 infection with respiratory symptoms .10 of them had pancreatitis of mild severity ,2 developed severe necrotizing pancreatitis.

Interpretation and conclusion: Based on our study we conclude that more than severe covid19infection mild covid 19infection with GI symptoms has a greater impact on the prognosis of patient with pancreatitis. However there is still insufficient evidence showing that covid 19 can cause TS or negatively impact prognosis. Additional major studies are needed to clarify relationship between these two entities

Keywords: Acute pancreatitis, Gastro intestinal symptoms, corona virus disease 2019

Introduction

Gastrointestinal symptoms are highly prevalent in corona virus disease ranging from 17.6 % to 53%.The proposed mechanism for GI symptoms involves SARS-COV2 virus binding to the host epithelial cell's ACE 2 receptor, commonly found in GI tract epithelial cells.

After defeating covid to an extent there is now an interest in understanding post covid sequele .“Acute pancreatitis ”is usually a self-limiting disease; however, severe form of the disease developed in 25 %of patients & it is associated with a mortality of up to 50 %. Available Scoring system’s aim is to stratify the severity of the T S, and this in turn guides the management with improving outcomes.

In this study we are tried understand if there was any impact of covid 19 infection on causation or

prognosis of pancreatitis .

PATHOGENESIS OF ACUTE PANCREATITIS

The initial phase is characterized by intra pancreatic digestive enzyme activation and acinar cell injury. Trypsin activation appears to be mediated by lysosomal hydrolases such as cathepsin B that become colocalized with digestive enzymes in intracellular organelles; it is currently believed that acinar cell injury is the consequence of trypsin activation.

The second phase of pancreatitis involves the activation, chemo attraction and sequestration of leukocytes and macrophages in the pancreas resulting in an enhanced intrapancreatic inflammatory reaction. Neutrophil depletion induced by prior administration of an anti neutrophil serum has been shown to reduce the severity of experimentally induced pancreatitis. There is also evidence to support the concept that neutrophils can activate trypsinogen.

Materials and Methods:

Source of data:

All patients diagnosed with acute pancreatitis who are hospitalized to the intensive care unit at the Ghandi Institute of Medical Sciences and Research Centre will be included.

A. Study design: Prospective Observational study

B. Study period 18 months February 2021 to August 2022

C. Place of study: Research Facility of the Ghandi Institute of Medical Sciences.

D. Sample Size: 30

E. Inclusion criteria:

- Patients with acute pancreatitis
- Prior history of covid19 infection
- People who are willing to provide informed consent

F. Exclusion Criteria:

- ◆ Chronic pancreatitis patient
- ◆ Patient with on going covid 19 infection
- ◆ Immuno compromised patients

G. Methodology:

Clinical and laboratory data of all consecutive patients with a primary diagnosis of TS during past 6 months with history of covid 19 infection was collected .Clinical and radiological grading was taken into consideration for finding severity.

SEVERITY OF ACUTE PANCREATITIS

Mild, Moderately severe, and severe

MILD ACUTE PANCREATITIS

- Without local complications or organ failure.
- Mostly associated with interstitial acute pancreatitis
- The disease is self-limited and subsides spontaneously

MODERATELY SEVERE ACUTE PANCREATITIS

- Transient organ failure (resolves in <48 h) or local or systemic complications in the absence of persistent organ failure.
- These patients may or may not have necrosis
- May develop local complication such as a fluid collection that requires a prolonged hospitalization greater than 1 week

SEVERE ACUTE PANCREATITIS

- Characterized by persistent organ failure (>48h).
- Organ failure can be single or multiple

Risk Factors for Severity

- Age > 60 years
- Obesity
- BMI > 30
- Comorbid disease (Charlson comorbidity Index)

Markers of Severity at Admission or Within 24hrs.

- SIRS defined by presence of 2 or more criteria
- Core temperature < 36°C or > 38°C
- Heart rate > 90 beats/min
- Respirations > 20/min or Pco₂ < 32 mmHg
- White blood cell count > 12,000/micL or < 4,000/micL or 10% bands
- APACHE II
- Hemoconcentration (hematocrit > 44%)
- Admission BUN (> 22 mg/dL)
- BISAP score

(B) BUN > 25 mg/dL

(I) impaired mental status

(S) SIRS > 2 of 4 present

(A) Age > 60 years

(P)Pleural effusion

- Organ failure(Modified Marshall 5 core)
- Cardiovascular: systolic BP < 90 mmHg, heart rate > 130 beats/min. Pulmonary: Pao₂ < 60 mmHg
- Renal: serum creatinine > 2.0 mg%

Markers of Severity during Hospitalization

- Persistent organ failure


STATISTICAL ANALYSIS:

Using SPSS V.20 for analysis, the obtained data will be entered into Microsoft Excel. The result will be expressed in the form of descriptive and inherently statistics.


If $p < 0.05$, it is said to be statistically significant

Results:

Out of 30 patient 4 patients had severe covid 19 infection requiring more than 10 days of hospitalization but though 1 of those 4 patient developed recurrent pancreatitis but severity score was less (clinical and radiological).




12 patients had mild covid 19 infection with respiratory symptoms. 10 of them had pancreatitis of mild severity, 2 developed severe necrotizing pancreatitis.



14 had mild covid 19 infection with minimal respiratory and more of GI symptoms .Out of these 14 ,10 developed severe acute necrotizing pancreatitis requiring ICU stay and prolonged hospital stay.

MILD PANCREATITIS VS SEVERITY OF COVID 19 PNEUMONIA



Discussion:

Acute pancreatitis as a manifestation of COVID-19 infection has been reported in isolated case reports and case series, however, a large, national study has yet to be published

One theory for pancreatitis development is through the binding of ACE2 receptors, which are present in pancreatic ductal, acinar, and islet cells In amdar et al. describe an increase in the incidence of TS due to un-identified causes in COVID-19 positive patients compared to COVID-19 negative patients, implicating Sars-CoV-2 to be a possible causative agent

Our study is not without limitations but our study does emphasis on need for larger studies to prove correlation of covid 19 infection and Acute pancreatitis.

Our study gives an idea on correlation of severity of pancreatitis and severity of covid19 infection.

Conclusion:

Despite the trend in recent literature of trying to establish or refute the role of SARS-CoV-2 in TS cases, currently, there is no sufficient evidence showing that COVID-19 can cause TS or negatively impact prognosis. Adherence to TS guidelines, namely diagnosis and etiological work-up, and careful monitoring of patients are of utmost importance to ensure the most adequate orientation and avoid convenience diagnosis. Based on our study we conclude that more than severe covid 19 infection mild covid 19 infection with GI symptoms has a greater impact on the prognosis of patient with pancreatitis. However there is still insufficient evidence showing that covid 19 can cause TS or negatively impact prognosis. Additional major studies are needed to clarify relationship between these two entities.

Acknowledgement

Author would like extend thanks to the Department of Zoology for carried out this research work, and Ghandhi Institute of Medical sciences.

Conflict of Interest: The author has no conflict of Interest.

Reference

1. Worldometer W. Coronavirus Update (live)-Worldometer.[cited25December 2020].Available from:

<https://www.worldometers.info/coronavirus/>

2. GuanWJ, NiZY, HuY, LiangWH, OuCQ, HeJX, LiuL, ShanH, LeiCL, HuiDSC, DuB, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020;**382**:1708–1720. [[PMCfree article](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* 2020;**158**:1831–1833. :e3. [[PMCfreearticle](#)] [[PubMed](#)] [[GoogleScholar](#)]
4. Bourgonje AR, Abdulle AE, Timens W, Hillebrands JL, Navis GJ, Gordijn SJ, Bolling MC, Dijkstra G, Voors AA, Osterhaus AD, van der Voort PH, Mulder DJ, van Goor H. Angiotensin- converting enzyme 2 (ACE2), SARS-CoV-2 and the pathophysiology of coronavirus disease 2019 (COVID-19) *J Pathol*. 2020;**251**:228–248. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
5. MaoR, QiuY, HeJS, TanJY, LiXH, LiangJ, ShenJ, ZhuLR, Chen Y, IacucciM, NgSC, Ghosh S, Chen MH. Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol*. 2020;**5**:667–678. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
6. Liu F, Long X, Zhang B, Zhang W, Chen X, Zhang Z. ACE2 Expression in Pancreas May Cause Pancreatic Damage After SARS-CoV-2 Infection. *Clin Gastroenterol Hepatol* 2020;**18**: 2128- 2130. :e2. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
7. Moher D, Liberati A, Tetzlaff J, Altman DG PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg*. 2010;**8**:336–341. [[PubMed](#)] [[Google Scholar](#)]
8. Lax SF, Skok K, Zechner P, Kessler HH, Kaufmann N, Koelblinger C, Vander K, Bargfrieder U, Trauner M. Pulmonary Arterial Thrombosis in COVID-19 With Fatal Outcome : Results From a Prospective, Single-Center, Clinicopathologic Case Series. *Ann Intern Med*. 2020;**173**:350–361. [[PMCfreearticle](#)] [[PubMed](#)] [[GoogleScholar](#)]
9. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, Tsiotos GG, Vege SS Acute Pancreatitis Classification Working Group. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut*. 2013;**62**:102–111. [[PubMed](#)] [[Google Scholar](#)]
10. Zhang H, Kang Z, Gong H, Xu D, Wang J, Li Z, Cui X, Xiao J, Zhan J, Meng T, Zhou W, Liu J, Xu H. Digestive system is a potential route of COVID-19: an analysis of single-cell coexpression pattern of key proteins in viral entry process. *Gut*. 2020;**69**:1010–1018. [[Google Scholar](#)]