International Journal for Multidisciplinary Research (IJFMR)
E-ISSN: 2582-2160 ● Website: www.ijfmr.com ● Email: editor@ijfmr.com

Neuropsychological Insights into Sensory-Based Intervention: The Role of Tactile and Vestibular Activities in Emotional and Motor Regulation during Early Milestones

Ms. Radha Navnit Jajal[1], Ms. Badri Nirav Patel[2], Ms. Khushi Prasad Cholkar[3]

[1]Clinical Neuropsychologist, Psychology, Healvibe Clinic
[2]Psychologist, Psychology, Healvibe Clinic
[3]Student, Psychology, Healvibe Clinic

ABSTRACT:
This study explored the potential of basic and organized sensory tasks to enhance tactile and vestibular control in Veda, a four-year-old child with mild oppositional defiant disorder (ODD) and learning disability. Children with such issues tend to have difficulties processing sensory information, body stability, and fine motor skills. These issues may influence learning, behavior, and emotional control. The current study addressed these issues through systematic sensory-motor pre-, post-, and intervention tests. These included Velcro pick-up-drop tasks, walking barefoot on various textured mats, walking and balancing on wooden beams, and board tasks to improve Veda's tactile tolerance, motor control, and postural control.
Veda was repeatedly tested for her sense of touch and balance through frequent sensory activities. Every time she walked on differently textured mats or stood on a beam, her brain had to simultaneously determine the movement in her body and what she was feeling. This eventually helped her brain become more adept at integrating these sensations, a process referred to as sensorimotor integration. This theory posits that the brain continuously receives and organizes sensory information to facilitate action. The most important systems in this process are the vestibular system, which allows us to sense movement and maintain balance, and the proprioceptive system that processes information about muscles and joints. Any failure in either of these systems can result in children presenting atypical responses concerning hypersensitivity, fear of movement, or even repetitive behaviors such as hand flapping.
A study conducted by A. Jean Ayres reveals that the brain uses the vestibular system to determine whether to respond to a stimulus. (Guardado & Sergent, 2023) These activities enhanced the activity of the sensorimotor strip of the brain, which performs touch and movement by improving its integration with the orbitofrontal cortex, which processes emotions and decision-making. As these two areas began to communicate more easily, Veda became less avoidant and less receptive, and combined sensory-related interventions have the potential to enhance the manner in which the brain processes and organizes sensory information. The findings can lead to the creation of more effective interventions to help children who are easily overstimulated by sensory stimuli and have developmental problems.

KEYWORDS: Sensory processing; sensory-motor strip; tactile sensitivity and tolerance; vestibular integration; orbitofrontal cortex; emotional regulation; oppositional defiant disorder; neuropsychological development; balance and postural control; early childhood intervention

INTRODUCTION:

Sensory and motor impairments are common in children with minor behavioral and learning problems, including oppositional defiant disorder and learning disabilities. They can be hypersensitive to touch, avoid balancing activities, or show frustration when they encounter unexpected sensations. These are not just behavioral responses but are directly related to the neuropsychological functioning of the way the brain processes sensory input and/or coordinates motor behavior. For the sake of developing effective therapeutic interventions in the future, understanding this relationship neuropsychologically holds the highest priority.

The orbitofrontal cortex and sensory-motor strip are the main areas involved. One of the subdivisions of the frontal cortex is the orbitofrontal cortex, which modulates the affective states and reward-based decision-making. For example, when a child is exposed to an unfamiliar touch, such as bubble wrap, the orbitofrontal cortex differentiates between approach and avoidance responses and thus orchestrates the corresponding affective responses, such as fear or engagement. The structural or functional differences observed in this area are common in children with ODD and explain their increased reactivity to challenging tasks.

Moreover, the sensorimotor cortex has two main areas:

1. **Sensory cortex:**

The sensory cortex is the area where data obtained by the five senses, namely, tactile, olfactory, gustatory, visual, and auditory senses are collated and evaluated. Furthermore, the neural signals were grouped or segregated into conscious perceptions. Every sensory modality is then anatomically divided into primary and secondary cortical areas; the primary areas receive the first afferent input, whereas the secondary levels carry out higher-level processing of such inputs, integrating them into multimodal associative networks. (ScienceDirect, n.d.)

2. **Motor cortex:**

The motor cortex is the part of the human brain that regulates voluntary movement by transmitting corticospinal impulses to the skeletal muscles. Its other major roles include the initiation of movement, coordination of complex actions, and regulation of balance and posture. (ScienceDirect, n.d.)

In our study, we relied on new sensory interventions to achieve reliable activation of these brain regions. A pre-test was conducted, whereby the participants underwent an organized sensorimotor activity, followed by a post-test, where the change in performance and behavior was measured. Through the constant repetition of such activities, new neural memories can be formed, which can yield better results. Neural memory refers to the mechanism through which the brain memorializes and holds experiences by developing and strengthening neuronal networks. This encompasses neural plasticity, which helps in the flexibility of the brain and learning of new patterns of response. There is repetitive stimulation reinforcing a stable association between sensory cortices, which are the main receivers of external input, and frontal cortical areas, which are involved in emotional and behavioral regulation in children with Oppositional Defiant Disorder (ODD) or learning disabilities. (Queensland Brain Institute, n.d.)

In the investigation under consideration, all sensory activities act as training in a protocol that strengthens neural memory records, thus increasing postural stability, fine motor performance, and less disturbed affective reactions. Therefore, chronic sensory-motor involvement not only alleviates behavioral performance but also redesigns neurocognitive mechanisms that control the perception of the environment and reaction to it.

It also has several clinically relevant outcomes, such as:

1. Emotional regulation improves in settings that present new sensory challenges.
2. Improved motor coordination for tasks, including balance, writing, and walking.
3. Enhanced attention and willingness to learn in the classroom.

In addition, the findings of this study can assist healthcare professionals, teachers, and parents in developing play-based sensory interventions that are not only in accordance with current neuroscience but also stimulate the interests of a child.

PURPOSE OF THE STUDY:

The problem that this study aims to answer is the possibility of connecting pleasurable childhood experiences with the neuropsychological mechanisms that influence emotional, behavioral, and cognitive development.

In particular, it dwells upon the case of Veda, a four-year-old child with mild Oppositional Defiant Disorder (ODD), learning impairment, and speech and language development delays. The exercises involved walking on textured mats with bare feet, balancing on a beam, and pick-and-drop exercises. They assist the child in developing sensory tolerance, motor coordination, emotional control, and motivation to learn. They do not just stimulate the brain by routine exercise, but also enhance communication between the essential brain areas, mainly the sensorimotor and orbitofrontal cortex.

This study neuropsychologically tested the effects of repeated sensorimotor stimuli on neural integration and plasticity. Neural memory is based on the plasticity of synaptic connections between neurons, with repeated experiences reinforcing the connections between neurons. The neurons that are concerned with the repetition of a particular activity will become more likely to fire together, and they will strengthen their connection because of long-term potentiation. (National Library of Medicine, n.d.).

Over time, consistent sensory experiences promote myelination, which accelerates signal transmission, leading to improved emotional control, focus, and motor coordination. Such mechanisms aid the brain in developing stronger and more efficient linkages in the areas concerned with processing emotion, movement, and sensation. Communication between these parts of the brain may not be fully functional in children, such as Veda, who are emotionally reactive, avoidant, and face learning challenges. Individuals indulging in playful and goal-oriented sensory activities provide their brains with repeated feedback. This assists in aligning the perceptions and actions. This may have resulted in her being less emotionally defensive, more focused, and more confident with her movements. These reforms are important for improving students' learning and communication abilities.

In addition, this study is not only aimed at the personal development of Veda. This study aimed to prove the feasibility of early childhood intervention using sensory play as a neuropsychologically grounded therapeutic model. It also shows how routine sensory activities contribute to the formation of new brain routes and enhance developmental outcomes by noting behavioral changes, the development of coordination, and affective responses before and after the intervention. In doing so, it identifies the important interdependence between neuroscience and early childhood education. When therapies correspond to the processes of natural learning, including movement, sensation, and play, they can result in major changes in behavior and thinking processes. Finally, it assists us in realizing how sensorimotor processes can contribute to the development of physical and emotional well-being, communication, and self-regulation in young children with developmental difficulties.

RELEVANCE OF THE STUDY:

This study is meaningful because it adds to our knowledge of the effects of sensory-related interventions on improving the neuropsychological development of young children with learning and behavioral difficulties. It included simple but highly designed sensory tasks, including the pick-and-drop task, balance beam, and walking barefoot on different textures of surfaces. The generalizability of this experiment is not limited to a specific case because it introduces a sound framework that unites early childhood therapy and neuroscience in a wide-ranging and adaptable play-based approach. These findings are important for professionals dealing with children with similar problems, such as therapists, educators, and clinicians.

Another important aspect of this research is that it helps to develop a new area of neurodevelopmental rehabilitation: the process of stimulating the connection between the neurons of the brain through experience and repetition. Connecting the roles of tactile and vestibular activities in the functions of the sensory, motor, and orbitofrontal cortices provides evidence that particular sensory-motor exercises can contribute to better regulation of emotions, postural control, and effectiveness of tasks.

Finally, this research is relevant because it connects scientific knowledge with real-world practices. It invites specialists to consider sensory play as a method of neurodevelopment, relying on a specific stimulation of the neural network through which emotional and behavioral reactions can be restructured. This study provides an effective theoretical framework and intervention paradigm that can be easily implemented, which is why it can become a valuable asset to any person attempting to facilitate the comprehensive development of young children with learning, emotional, or sensory problems. The results indicate that with planned early interventions, with the sense of how the brain takes the input of sensory information, it can change considerably and permanently the development of a child, their communication, and well-being.

RESEARCH QUESTION:

How can repetitive, play-based sensory experiences improve neural integration between the sensory, motor, and emotional regulation systems in a four-year-old child with oppositional defiant disorder and learning disability?

LITERATURE REVIEW:

1. **Dunn's Model of Sensory Processing:** Dunn's model of sensory processing offers a theoretical basis for explaining the differences in how children perceive and react to sensory input. The model postulates that every individual has a definite neurological threshold that dictates the amount of stimulation required to produce a response, as well as behavioral strategies such as seeking, avoidance, or registration of stimuli. Applying this framework in the present study, the sensory variability of Veda was analyzed through structured activities, including balance exercises, textured walks, and pick-and-drop activities aimed at modulating sensory thresholds. (Alexia E Metz, 2019)
2. **Process Model of Emotional Regulation:** The process Model of Emotion Regulation is a theory developed by Gross, according to which people attempt to maintain affective states using several techniques, such as attentional deployment, cognitive reappraisal, and response modulation. Through sensory play, repeated exposure to such activities allows children to implement these strategies in a

safe and structured environment. Veda, who first showed increased emotional reactivity and avoidance, used these experiences to divert attention, recontextualize discomfort, and regulate impulsive responses. (GROSS, 2002)

3. **Hebbian Learning and predictive mirror neurons for action, sensations, and emotions:** Recent neuroscience studies have indicated that the brain enhances the association among neurons when certain experiences are repeated. Keysers and Gassola argue that as children repeatedly associate what they feel, see, and do, their brains make more connections between the sensory, motor, and emotional regions. Their predictive mirror neuron theory explains that the brain does not simply mirror what others do but also predicts and reacts better to sensory input. (Christian Keysers, 2014)

CASE STUDY:

Veda, a 4-year-old girl, was presented to the clinic by her parents because of concerns regarding developmental delays, behavioral issues, and problems with sensory regulation impacting her everyday functioning. Her parents claimed that she always demonstrated defiant behavior, frequent tantrums, and a poor understanding and ability to follow easy instructions. These difficulties were also noted at home and in early learning settings, where she experienced difficulties with attention, socialization, and group activities.

Clinically, she was alert and cooperative but faced difficulty in sustaining attention for a long duration. She was distracted by environmental stimuli and had to be redirected frequently during the course of the tasks. It was observed that she had mild muscular weakness in her upper limbs and a lack of fine motor coordination, especially regarding grip and precision. She regularly used compensatory postures to maintain balance, such as a wide stance or occasional toe walking.

Sensory assessment, performed using the VORT Sensory Checklist, showed that she was hypersensitive to touch and proprioception. She prefers not to touch different textures of objects. She also exhibited repetitive hand movements, which were probably a form of self-regulation. These results demonstrate that she experienced sensory modulation issues that influenced her attention and adaptive behavior.

A medical examination, consistent with DSM-5, revealed that Veda had characteristics of Specific Learning Disability and Oppositional Defiant Disorder. These diagnoses present themselves in the form of persistent attention problems, an inability to complete tasks, and an overall academic deficiency that is particularly impaired in challenging cognitive tasks. Defiant reactions, resistance to instruction, and difficulties in emotional control usually follow these symptoms.

After the intervention period, which used systematic sensorimotor and multisensory treatment, Veda showed remarkable improvements in her involvement and performance. The play-based therapeutic processes focused on tactile and vestibular processing, producing higher tolerance to sensory inputs, better fine and gross motor coordination, and emotional regulation. The difficulty of the vestibular task gradually increased at the end of the post-test phase, during which she could perform the tests with confidence and less avoidance. Her attention and speed of completion of tasks improved, and her tantrums reduced.

Altogether, this case is an example of a child with neurodevelopmental and sensory integration impairments who received significant value from intensely designed and play-based interventions. Her development demonstrates the importance of early diagnosis and an individual treatment plan for developing cognitive and emotional growth.

METHODOLOGY:

The following research with Veda employed a straightforward three-phase structure, namely, pre-test, intervention, and post-test, to investigate the role of structured sensory activities on the functions of tactile and vestibular processing, emotional regulation, and motor coordination. Every stage aimed to gather both quantitative and qualitative information, all in the framework of a play-based and child-centered approach that would encourage elements of intrinsic motivation and emotional safety.

In the pre-test phase, Veda underwent an intensive baseline test using the VORT Sensory Checklist. This test helped us determine her sensory modulation patterns in the tactile, vestibular, proprioceptive, visual, and auditory systems. The findings provided a full account of her strengths and weaknesses.

Nonetheless, our research primarily focused on the tactile and vestibular areas. The obtained profile formed the foundation for individualized intervention planning. Fine motor coordination, sensory registration, and the ability to control attention were measured using additional observational and task-based tests, including the Velcro pick-and-drop task and balance beam and board tasks. These activities helped identify specific challenges in daily activities. She showed moderately high tactile defensiveness and indecisiveness in balance activities, indicating a tendency for sensory overload and high emotional reactivity to unknown physical challenges.

Veda participated in a program of sensorimotor sessions throughout the course of the intervention. These sessions were organized over time. The program aimed to enhance sensory integration, self-control, and coordination. We employed barefoot textured walks and graded activities involving the use of a vestibular, such as balancing on a board, to assist in improving postural control and spatial awareness.

Intervention became more challenging and demanding as the program progressed. This test assessed the balance, attention, and emotional adaptability of the Veda. She became more confident and controlled, even with the increased demand. Regular support and encouragement of the systematic intervention led to her addressing each task with no hesitation and reduced her fatigue. It also assisted in reducing her anxiety during performance.

In the post-test phase, we repeated the same sensory and behavioral measures to analyze the progress. Changes in sensory responsiveness, attention, and affective control after the intervention were noted again using the VORT Sensory Checklist. There was an improvement in performance on the Velcro pick-and-drop task and balance beam, with increased accuracy, sustained attention, and fluency in motor sequencing. Veda developed better emotional management and self-awareness as her emotional responses changed from avoidance and irritability to curiosity and persistence.

Ethical guidelines were followed throughout the study. Veda's parents provided informed consent to initiate the intervention and discuss its purpose, procedures, and anticipated benefits. Each session was conducted in a secure and safe environment and was controlled with the help of trained professionals in child development and neuropsychology. We also ensured that no respondent was physically or emotionally uncomfortable. No records were disclosed, and no report on the subject was provided to reveal her identity.

TASKS	TYPE	BRAIN REGIONS INVOLVED	KEY FUNCTIONS
Velcro pick up drop	Tactile	Primary somatosensory cortex Primary motor cortex	Processes tactile information like touch and pain.

		Parietal lobe	Planning and execution Spatial orientation
Textured mat walking	Tactile	Somatosensory cortex Supplementary motor area Posterior parietal cortex	Emotional processing Movement planning Eye-hand coordination
Balance Beam	Vestibular	Basal ganglia Vestibular cortex Brainstem (vestibulospinal tract)	Postural control Maintaining balance Maintaining head position and stability
Balance Board	Vestibular	Vestibular nuclei Cerebellum (flocculonodular lobe) Parietal lobe	Sensory integration (visual input and proprioception) Processes signals from the inner ear's vestibular system to maintain equilibrium Spatial navigation

RESULTS:

1. **Velcro pick up-drop task:**

Pre-test			Post-test		
Number of trials	Percentage of Hits	Percentage of Misses	Number of trials	Percentage of Hits	Percentage of Misses
1	67	33	1	67	33
2	67	33	2	83	17
3	67	33	3	100	0
4	83	17	4	100	0
5	100	0	5	100	0
6	100	0	6	100	0

2. **Textured mat walking task (intervention):**

Number of trials	Percentage of Hits	Percentage of Misses
1	80	20
2	90	10

3. **Balance beam task (pre-test):**

Number of trials	Percentage of Hits	Percentage of Misses
1	40	60
2	90	10

4. Balance board task (post-test):

Number of trials	Percentage of Hits	Percentage of Misses
1	50	50
2	80	20

INTERPRETATION:

1. VELCRO PICK-AND-DROP TASK





The outcome of the pre- and post-test of the Velcro pick-and-drop activity is presented in the graphs below. The pre-test graph indicates a lack of consistency after 6 trials. Hit rates were found to be between 65 to 100%, and miss rates were 15 to 35% range. These differences indicate that she had slower sensory processing, her motor control was unstable, and she had less attention to tactile feedback. She also had poor visual-spatial attention, thus failing in coordination and accuracy.

Conversely, the post-test graph illustrates a noticeable and consistent increase in accuracy as well as precision. The proportion of the hits increases gradually to 100% in the third trial, whereas the misses decrease rapidly to 0-20%. Such an improvement translates to enhanced tactile responses, motor planning, and increased hand-eye coordination in Veda. Although the post-test tasks were more demanding in terms of accuracy and control, she performed them more uniformly, with reduced effort. This development indicates that she organized sensory play intervention improved her sensory-motor integration, tactile sensitivity, and her ability to focus and have executive control when performing tasks.

2. TEXTURED MAT WALK TASK



There is a large improvement in the graph of tactile awareness between round 1 and round 2. Her hit rate in round 1 was 80% and her miss rate was 20%. She exhibited fatigue, hesitation, and avoidance, flicking her hands and losing attention in the middle of the task. This led to an increased short-sightedness and made the task hard to follow, demonstrating difficulty in maintaining tactile focus.

During round 2, she was more efficient with 90% of hits and 10% of misses. She was less fatigued, less avoidant, and more engaged and confident of the tactile sensations. This resulted in increased sensory processing and awareness, as seen in improved consistency in performance. These improvements prove that frequent practice and a formal intervention enabled her to become more open to the tactile stimuli and incorporate sensory feedback into her behavioural patterns.

3. BALANCE BEAM AND BOARD TASK





The table shows the pre- and post-testing outcomes of the balance beam and board task. Hits in round 1 were 40%, and misses were 60%, which means that there was a challenge in associating body movements with sensory information, and a lack of control and precision. During round 2, the hits went up to 90%, and the misses dropped to 10%, which shows that with repetition, she was able to take a steady position and was able to do the task better.

It is supported by post-test data showing that the sensory-motor intervention affected her performance. During round 1, both the hit and miss rates were 50%, indicating that balance and control remained a problem. When the task was complicated, round 2 produced an 80% hit rate and a 20% miss rate, which is an indication of improved control, coordination, and concentration than the pre-test.

Furthermore, the results also imply that she could maintain her posture, balance on the board more accurately without any fear. It shows that structured sensory intervention was successful in supporting her overall motor development and adaptive responses.

LIMITATIONS:

Although the current research has some drawbacks, it also highlights the excellence and complexity of the research design and its relevance to neurodevelopmental psychology.

1. **Single-participant focus as a deep case exploration:**
   The study design focused on one participant, but this configuration allowed making a high-resolution study of behavioural and neuropsychological changes over time. This intensive, personalized methodology has been able to record finer nuances in tactile thresholds, motor control, and emotion regulation that would otherwise be dampened down in population-size research. The accuracy makes the results useful in the creation of individual intervention models of pediatric neuropsychology.
2. **Limited multidimensional outcome measures:**
   This investigation centred largely on tactile and vestibular reactions. Inclusion of tasks measuring attention span, emotional regulation, or executive functioning could provide a thorough account of the process of multisensory integration.
3. **Short-term, but quickly changing neurobehavioral:**
   The fact that the apparent progress was made in a short period of time despite the brief duration of intervention highlights the vulnerability of the developing brain to the received sensory input. This

immediate reaction offers evidence at early stages of development in a child with developmental difficulties, which makes the results important in a clinical and theoretical manner.

RECOMMENDATIONS:

1. Further research must enroll bigger cohorts that are likely to represent the rest of the population with varying age, gender, ethnicity, socioeconomic status, and other relevant demographic variables, in a bid to improve external validity. They will be required to carry out comparative analyses on children with heterogeneous sensory profiles (hypersensitivity, hyposensitivity, or mixed modulation patterns) to determine the existence of a difference in the efficacy of intervention across sensory subtypes. Causal inference would be further enhanced by randomized controlled trials that, in the future, could decouple the particular contribution of sensory-motor interventions.
2. The sessions should also incorporate predetermined relaxation measures, including deep-pressure tactile stimulation, directed diaphragmatic breathing, and rhythmic movement protocols, so that there could be a unification of sensory input and affective regulation. Such practices could improve emotional regulation abilities in children by encouraging stability of the limbic-prefrontal circuitry, especially that of the orbitofrontal cortex.
3. Prolonging intervention phases along with a systemic long-term follow-up assessment will be necessary to establish whether the described behavioral and motor improvements are maintained, transferred to real-life situations, and relevant to other areas of development, such as attention control, school preparation, and social interaction.
4. Finally, include outcomes of feasibility and acceptability, such as parent and therapist feedback, child involvement, and negative events. In case the findings are promising, run pragmatic trials and cost-efficiency evaluations to determine how it can be scaled or create open-access intervention materials (manuals, short training videos) to enable clinicians and educators to run evidence-based versions of the protocol and still achieve an outcome. These will transform the protocol into an informative presentation of a case with stringent clinically applicable evidence on the sensory-motor interventions in young children with ODD and learning issues.

CONCLUSION:

The current study provides solid empirical data, in which systematic sensorimotor and multisensory interventions instigate significant improvements in fine motor control, sensory discrimination, balance, and attentional stability among developmentally delayed children. The results obtained can support the idea of experience-related neuroplasticity, according to which extended sensory and motor stimulation can promote cortical reorganization and thalamocortical coordination.

The reduction in fatigue and increase in attention during intervention indicated the enhancement of neural efficiency and regulation. These effects provide neuropsychological evidence in support of the theory that early multisensory exposure is capable of recalibrating atypical sensory thresholds and, hence, enhancing self-regulation, confidence, and readiness to learn.

As a whole, these studies span the divide between neuropsychological theory and early intervention practice, and show that through targeted, play-based sensory interventions, meaningful change can occur in a short space of time. Although further research is required, the findings of the research have a high likelihood of confirming the integration of neuroscience-informed sensory play in early childhood therapy and education, to enforce balanced emotional, sensory, and cognitive development. (Dave Neale, 2018)

REFERENCES:

1. Duch, W., Bekkering, H., & Neggers, B. Sensorimotor integration. Integration The Vlsi Journal, 1-15.
2. Guardado, K. E., & Sergent, S. R. (2023, July 31). *Sensory Integration*. StatPearls - NCBI Bookshelf. https://www.ncbi.nlm.nih.gov/books/NBK559155/
3. Motor Cortex - an overview | ScienceDirect Topics. (n.d.). www.sciencedirect.com. https://www.sciencedirect.com/topics/medicine-and-dentistry/motor-cortex
4. Sensory Cortex - an overview | ScienceDirect Topics. (n.d.). www.sciencedirect.com. https://www.sciencedirect.com/topics/medicine-and-dentistry/sensory-cortex
5. Queensland Brain Institute. (2016, December 2). How are memories formed? uq.edu.au. https://qbi.uq.edu.au/memory/how-are-memories-formed
6. Purves, D., Augustine, G. J., Fitzpatrick, D., Katz, L. C., LaMantia, A.-S., McNamara, J. O., & Williams, S. M. (2001). Long-Term Synaptic Potentiation. Neuroscience. 2nd Edition, 2. https://www.ncbi.nlm.nih.gov/books/NBK10878/
7. Metz, A. E., Boling, D., DeVore, A., Holladay, H., Liao, J. F., & Vlutch, K. V. (2019). Dunn's Model of Sensory Processing: An Investigation of the Axes of the Four-Quadrant Model in Healthy Adults. Brain Sciences, 9(2), 35. https://doi.org/10.3390/brainsci9020035
8. Gross, J. J. (2002). Emotion regulation: Affective, cognitive, and Social Consequences. Psychophysiology, *39*(3),281–291. https://doi.org/10.1017/s0048577201393198
9. Keysers, C., & Gazzola, V. (2014). Hebbian learning and predictive mirror neurons for actions, sensations and emotions. Philosophical Transactions of the Royal Society B: Biological Sciences, 369(1644), 20130175. https://doi.org/10.1098/rstb.2013.0175
10. Neale, D., Clackson, K., Georgieva, S., Dedetas, H., Scarpate, M., Wass, S., & Leong, V. (2018). Toward a Neuroscientific Understanding of Play: A Dimensional Coding Framework for Analyzing Infant–Adult Play Patterns. *Frontiers in Psychology*, *9*. https://doi.org/10.3389/fpsyg.2018.00273