International Journal for Multidisciplinary Research (IJFMR)
E-ISSN: 2582-2160 ● Website: www.ijfmr.com ● Email: editor@ijfmr.com

# Return to Sport Criteria Following ACL Reconstruction: Current Evidence and Physiotherapy Perspectives – A Narrative Review

**Dr. Amaresh Mohan[1], Dr. Bhupesh Goyal[2]**

[1]Research Scholar, Dept. of Physiotherapy, Vivekananda Global University, Jaipur, India.
[2]Professor & HOD, Dept. of Physiotherapy, Vivekananda Global University, Jaipur, India.

**Abstract**

**Background:** Anterior cruciate ligament (ACL) injury is one of the most prevalent sports-related knee injuries and is frequently associated with functional instability, reduced athletic performance, and prolonged absence from sports participation. Although ACL reconstruction (ACLR) is widely performed to restore knee stability, determining the optimal criteria for return to sport (RTS) remains a significant clinical challenge. Traditionally, RTS decisions have been based on postoperative timelines; however, recent evidence suggests that objective functional and performance-based assessments provide a safer and more reliable approach.

**Objective:** The purpose of this narrative review is to examine current evidence regarding return-to-sport criteria following ACL reconstruction and to highlight the role of physiotherapy in assessing readiness for sports participation.

**Methods:** A narrative review of the literature was conducted using published articles from peer-reviewed journals focusing on ACL reconstruction, rehabilitation, and return-to-sport assessment. Relevant studies, systematic reviews, clinical guidelines, and expert recommendations published between 2020 and 2026 were reviewed. Key areas explored included muscle strength assessment, functional performance testing, balance and proprioception evaluation, movement quality analysis, return-to-sport timing, and physiotherapy-based decision-making.

**Results:** Current evidence indicates that return-to-sport decisions should not rely solely on time elapsed since surgery. Physiotherapy-based RTS assessment involves a comprehensive evaluation of quadriceps and hamstring strength, limb symmetry index, hop performance tests, dynamic balance, neuromuscular control, and movement quality. Studies consistently demonstrate that athletes who achieve objective functional criteria before returning to sport exhibit lower reinjury rates and better long-term outcomes. Criterion-based rehabilitation programs emphasizing progressive strengthening, functional training, and sport-specific activities are associated with improved readiness for athletic participation.

**Conclusion:** Return-to-sport following ACL reconstruction should be guided by objective physiotherapy assessments rather than time-based criteria alone. Comprehensive evaluation of muscle strength, functional performance, balance, proprioception, and movement quality is essential to ensure safe and successful return to sports. Physiotherapists play a critical role in identifying residual deficits, monitoring rehabilitation progress, and facilitating evidence-based return-to-sport decision-making. Adoption of

standardized criterion-based RTS protocols may reduce reinjury risk and optimize functional outcomes following ACL reconstruction.

**Keywords:** Anterior Cruciate Ligament Reconstruction; Return to Sport; Physiotherapy Rehabilitation; Functional Performance; Muscle Strength; Hop Tests; Limb Symmetry Index; Sports Rehabilitation.

## Introduction

Anterior cruciate ligament (ACL) injuries are among the most common and clinically significant injuries affecting the knee joint, particularly in athletes and physically active individuals. The ACL is a primary stabilizing structure of the knee that prevents excessive anterior translation and rotational movement of the tibia relative to the femur. Injuries to the ACL frequently occur during sports involving sudden deceleration, pivoting, cutting, jumping, and landing maneuvers, such as football, soccer, basketball, volleyball, and rugby. The incidence of ACL injuries has increased globally due to growing participation in competitive and recreational sports, making ACL reconstruction (ACLR) one of the most commonly performed orthopedic surgical procedures.

The consequences of ACL injury extend beyond mechanical instability of the knee joint. Individuals often experience pain, recurrent episodes of giving way, reduced functional capacity, diminished athletic performance, and limitations in activities of daily living. Furthermore, untreated or inadequately rehabilitated ACL injuries may increase the risk of secondary meniscal damage, cartilage degeneration, and early-onset osteoarthritis. As a result, ACL reconstruction is frequently recommended for individuals wishing to return to high-demand physical activities and sports participation. The primary goals of ACL reconstruction and postoperative rehabilitation are to restore knee stability, regain functional performance, prevent reinjury, and facilitate a safe return to sport (RTS).

Return to sport is considered one of the most important outcomes following ACL reconstruction. For many athletes, successful return to their preinjury level of competition represents the ultimate measure of rehabilitation success. However, determining when an individual is ready to safely resume sporting activities remains a complex and challenging clinical decision. Historically, RTS decisions were largely based on the amount of time elapsed since surgery, with many athletes being cleared for sports participation between six and twelve months postoperatively. Although time-based criteria remain commonly used, contemporary evidence suggests that postoperative duration alone is insufficient for determining readiness to return to sport.

Research has demonstrated that many athletes continue to exhibit significant deficits in muscle strength, neuromuscular control, balance, movement quality, and functional performance even after completing standard rehabilitation programs. Returning to sports before achieving adequate physical recovery may increase the likelihood of graft failure, contralateral ACL injury, and recurrent knee instability. Studies have reported that athletes who return to high-level sports without meeting objective functional criteria face a substantially higher risk of sustaining a second ACL injury. Consequently, clinicians and researchers have increasingly emphasized the importance of criterion-based rehabilitation and return-to-sport assessment.

Physiotherapy plays a central role throughout the rehabilitation process following ACL reconstruction. Modern rehabilitation protocols focus not only on restoring joint range of motion and muscle strength but also on improving proprioception, dynamic stability, neuromuscular coordination, movement efficiency, and sport-specific performance. Physiotherapists are responsible for monitoring recovery progression,

identifying residual impairments, and implementing evidence-based interventions to optimize functional outcomes. Through systematic assessment and individualized rehabilitation planning, physiotherapists contribute significantly to reducing reinjury risk and enhancing long-term athletic performance.

Several objective measures have been proposed to evaluate readiness for return to sport following ACL reconstruction. These include assessments of quadriceps and hamstring muscle strength, limb symmetry index (LSI), single-leg hop tests, triple-hop tests, crossover hop tests, dynamic balance evaluations, and movement quality analyses. Functional performance testing has become an essential component of RTS decision-making because it provides valuable information regarding lower-limb strength, power, coordination, and symmetry. In addition, advancements in biomechanical assessment and motion analysis have enabled clinicians to identify subtle movement abnormalities that may not be apparent during routine clinical examinations.

Despite substantial progress in rehabilitation science, considerable variability continues to exist regarding RTS criteria and clearance protocols. Different rehabilitation centers and sports medicine practitioners often employ diverse assessment methods and cutoff values, leading to inconsistencies in clinical practice. Currently, there is no universally accepted return-to-sport framework that adequately addresses all aspects of physical recovery following ACL reconstruction. Therefore, the development and implementation of evidence-based RTS criteria remain a priority within sports physiotherapy and orthopedic rehabilitation.

The purpose of this narrative review is to examine the current evidence regarding return-to-sport criteria following ACL reconstruction from a physiotherapy perspective. Particular emphasis is placed on muscle strength assessment, functional performance testing, balance and proprioceptive evaluation, movement quality analysis, return-to-sport timing, and clinical decision-making. By synthesizing recent literature, this review aims to provide physiotherapists, rehabilitation professionals, and sports medicine practitioners with evidence-based recommendations to support safe and effective return-to-sport decisions following ACL reconstruction.

## ACL Reconstruction and Rehabilitation

### ACL Anatomy and Function:

The anterior cruciate ligament (ACL) is one of the primary stabilizing ligaments of the knee joint. It originates from the posteromedial aspect of the lateral femoral condyle and inserts into the anterior intercondylar region of the tibia. The ACL consists of two functional bundles: the anteromedial bundle and the posterolateral bundle. These bundles work together to provide stability throughout the knee's range of motion. The primary function of the ACL is to prevent excessive anterior translation of the tibia relative to the femur while also contributing to rotational stability of the knee.


The ACL plays a critical role during dynamic activities such as running, jumping, cutting, pivoting, and landing. Injury to this ligament results in altered knee biomechanics, impaired neuromuscular control, and functional instability, significantly affecting athletic performance and daily activities.

### Mechanism of ACL Injury:

ACL injuries may occur through contact or non-contact mechanisms. Non-contact injuries account for approximately 70–80% of ACL tears and commonly occur during rapid deceleration, sudden changes in direction, pivoting movements, or awkward landings from a jump. Excessive knee valgus, internal tibial rotation, and insufficient neuromuscular control are recognized risk factors for ACL injury.

Contact injuries occur when an external force is applied directly to the knee, such as during collisions in

football or rugby. Regardless of the mechanism, ACL rupture often results in immediate pain, swelling, joint instability, and inability to continue sporting activities.




**ACL Reconstruction:**

Anterior cruciate ligament reconstruction is considered the gold-standard surgical treatment for individuals seeking to return to physically demanding activities and sports. The procedure involves replacing the torn ACL with a graft to restore mechanical stability and knee function.

**Hamstring Tendon Autograft:**

Hamstring tendon grafts, typically harvested from the semitendinosus and gracilis tendons, are among the most commonly used graft options. Advantages include reduced anterior knee pain, smaller surgical incisions, and lower donor-site morbidity. However, postoperative hamstring weakness may occur during the early stages of rehabilitation.

**Bone–Patellar Tendon–Bone (BPTB) Autograft:**

The BPTB graft consists of the central third of the patellar tendon along with bone plugs from the patella and tibial tuberosity. This graft provides excellent fixation strength and has historically been considered the benchmark for ACL reconstruction. Potential disadvantages include anterior knee pain, patellar tendon discomfort, and difficulty kneeling.

**Quadriceps Tendon Autograft:**

The quadriceps tendon graft has gained popularity due to its large graft size and favorable biomechanical properties. Studies suggest comparable outcomes to hamstring and patellar tendon grafts, with lower don-

or-site morbidity in selected patients.
The choice of graft depends on patient characteristics, sporting demands, surgeon preference, and rehabilitation goals.

**Physiotherapy Rehabilitation Following ACL Reconstruction:**
Rehabilitation is a critical component of successful recovery following ACL reconstruction. The overall objective is to restore knee function, optimize physical performance, minimize complications, and facilitate a safe return to sport. Contemporary rehabilitation programs are criterion-based and progress according to functional milestones rather than fixed timelines.

**Phase I: Immediate Postoperative Phase (0–6 Weeks):**
The primary goals during this phase include controlling pain and swelling, protecting the graft, restoring knee range of motion, and re-establishing quadriceps activation.
Key interventions include:

- Cryotherapy and compression for edema control
- Patellar mobilization exercises
- Passive and active-assisted range of motion exercises
- Quadriceps setting exercises
- Straight-leg raises
- Weight-bearing progression as tolerated
- Gait training with assistive devices

Achieving full knee extension and minimizing quadriceps inhibition are critical priorities during this phase.

**Phase II: Early Strengthening Phase (6–12 Weeks):**
As pain and swelling decrease, rehabilitation progresses toward restoring muscular strength and neuromuscular control.
Major interventions include:

- Closed kinetic chain strengthening exercises
- Mini-squats and wall slides
- Leg press exercises
- Step-up and step-down training
- Static and dynamic balance exercises
- Core stabilization training
- Stationary cycling

The focus is on improving lower-limb strength while maintaining proper movement patterns and joint stability.

**Phase III: Advanced Strengthening and Neuromuscular Training (3–6 Months):**
This phase emphasizes development of muscular power, endurance, proprioception, and dynamic stability.
Common interventions include:

- Progressive resistance training
- Single-leg strengthening exercises
- Advanced balance and proprioceptive activities
- Agility ladder drills
- Plyometric exercises

- Lateral movement training
- Running progression programs

Physiotherapists carefully monitor movement quality to identify compensatory strategies and asymmetrical loading patterns.

**Phase IV: Return-to-Sport Training Phase (6–12 Months):**

The final phase prepares the athlete for sport-specific demands and return-to-sport testing.

Interventions include:

- Sprint training
- Cutting and pivoting drills
- Jump-landing mechanics training
- Sport-specific movement patterns
- Reactive agility exercises
- Functional performance testing
- Endurance conditioning

Emphasis is placed on achieving symmetrical strength, optimal movement quality, and sport-specific readiness.

**Importance of Criterion-Based Rehabilitation:**

Modern rehabilitation protocols advocate criterion-based progression rather than relying solely on time since surgery. Functional milestones such as restoration of range of motion, quadriceps strength recovery, balance performance, and successful completion of hop tests are considered essential indicators of readiness for advancement.

Criterion-based rehabilitation enables clinicians to individualize treatment according to patient progress and reduces the risk of premature return to sport. This approach is associated with improved functional outcomes, greater confidence in knee performance, and lower reinjury rates.

Successful ACL rehabilitation requires close collaboration between orthopedic surgeons, physiotherapists, strength and conditioning specialists, and athletes. Through comprehensive assessment and progressive exercise prescription, physiotherapy plays a pivotal role in restoring athletic performance and ensuring a safe return to sporting activities following ACL reconstruction.

**Return-to-Sport (RTS): Concept and Importance**

**Concept of Return-to-Sport:**

Return to sport (RTS) is a critical outcome measure following anterior cruciate ligament (ACL) reconstruction and rehabilitation. For athletes and physically active individuals, the ultimate objective of treatment is not only restoration of knee stability but also the ability to safely resume sporting activities at a level comparable to preinjury performance. RTS represents the culmination of the rehabilitation process and serves as an indicator of successful functional recovery.

Historically, return-to-sport decisions were primarily based on the amount of time elapsed since surgery. Athletes were often cleared to return to training and competition after a predetermined postoperative period, commonly between six and twelve months. However, contemporary research has demonstrated that recovery following ACL reconstruction is highly individualized, and athletes may exhibit significant deficits despite reaching traditional time-based milestones. Consequently, the concept of RTS has evolved

from a simple time-based decision to a comprehensive assessment of physical readiness and functional competence.

Modern sports rehabilitation emphasizes a criterion-based approach in which athletes must achieve specific functional, strength, and performance benchmarks before returning to sport. This approach aims to ensure that the reconstructed knee can tolerate the physical demands of athletic participation while minimizing the risk of reinjury.

**Stages of Return-to-Sport:**

Return to sport is not a single event but rather a progressive continuum consisting of multiple stages. The widely accepted model proposed by sports medicine experts includes three levels of return:

**Return to Participation:**

This stage involves the athlete resuming modified physical activities or team training without fully participating in competition. Participation may include non-contact drills, controlled practice sessions, conditioning exercises, and supervised sport-specific activities. At this stage, the athlete is physically active but has not yet achieved full competitive readiness.

**Return to Sport:**

Return to sport refers to the athlete resuming their chosen sport but not necessarily at their previous level of performance. Athletes may participate in competition while continuing rehabilitation and performance enhancement programs. Functional limitations may still exist, although participation is generally considered safe.

**Return to Performance:**

Return to performance represents the highest level of recovery. At this stage, the athlete has regained preinjury performance levels or achieved an even higher level of athletic capability. Physical performance, strength, agility, endurance, and sport-specific skills are fully restored, enabling unrestricted competition. Understanding these stages allows physiotherapists to establish realistic rehabilitation goals and monitor progress effectively.

**Importance of Return-to-Sport Assessment:**

The decision to return an athlete to sport is one of the most important responsibilities of sports rehabilitation professionals. Premature return to high-demand activities can result in serious consequences, including graft rupture, contralateral ACL injury, meniscal damage, and prolonged disability.

Comprehensive RTS assessment helps clinicians determine whether rehabilitation goals have been achieved and whether the athlete possesses the physical capabilities necessary for safe participation. Objective assessment also reduces reliance on subjective judgments and enhances consistency in clinical decision-making.

Physiotherapists play a central role in RTS evaluation through assessment of muscle strength, functional performance, balance, movement quality, and sport-specific abilities. These assessments provide valuable information regarding residual deficits that may increase injury risk.

**Risks Associated with Premature Return to Sport:**

Although many athletes are eager to resume competition as quickly as possible, returning to sport before adequate recovery may significantly increase reinjury risk.

Several factors contribute to elevated injury risk following ACL reconstruction:

- Persistent quadriceps weakness
- Reduced hamstring strength
- Impaired dynamic balance
- Poor neuromuscular control
- Altered movement mechanics
- Incomplete functional recovery
- Asymmetrical lower-limb loading

Athletes who return to sport without meeting objective RTS criteria often demonstrate compensatory movement patterns that place excessive stress on the reconstructed ligament and surrounding structures. Reinjury can result in additional surgery, prolonged rehabilitation, and reduced long-term athletic participation.

**Reinjury Following ACL Reconstruction**

A second ACL injury remains one of the most significant concerns after reconstruction. Reinjury may involve rupture of the reconstructed graft or injury to the contralateral ACL. Young athletes participating in cutting and pivoting sports appear to be particularly vulnerable.

Research indicates that reinjury rates are substantially higher among athletes who fail to achieve adequate strength symmetry and functional performance before returning to competition. Consequently, clinicians increasingly advocate comprehensive RTS testing batteries that evaluate multiple aspects of recovery rather than relying solely on surgical timelines.

**Physiotherapy Perspective on RTS Decision-Making:**

From a physiotherapy perspective, RTS decision-making should involve a multidimensional assessment process. Successful RTS requires more than restoration of passive knee stability; it demands optimal integration of strength, coordination, balance, endurance, agility, and movement efficiency.

Physiotherapists assess several domains during RTS evaluation, including:

- Knee range of motion
- Quadriceps and hamstring strength
- Limb symmetry index
- Functional hop performance
- Dynamic balance
- Proprioceptive function
- Agility and sport-specific skills
- Movement quality during jumping and landing

Objective testing allows clinicians to identify residual impairments and develop targeted interventions before sports participation is resumed.

**Need for Evidence-Based RTS Criteria:**

Despite significant advances in rehabilitation science, no universally accepted RTS protocol currently

exists. Variability in testing methods, performance thresholds, and rehabilitation practices continues to challenge clinicians. Therefore, evidence-based RTS criteria are essential to improve consistency and optimize patient outcomes.

Current literature supports the use of criterion-based rehabilitation and objective functional testing as key components of RTS decision-making. Such approaches facilitate individualized progression, reduce reinjury risk, and enhance long-term athletic performance.

In summary, return to sport represents a critical milestone following ACL reconstruction and requires careful evaluation of physical readiness. Physiotherapists play an indispensable role in assessing functional recovery and ensuring that athletes return to sports safely and effectively. A structured, evidence-based RTS framework remains fundamental to achieving successful rehabilitation outcomes.

**Physiotherapy-Based Return-to-Sport Criteria Following ACL Reconstruction:**

Determining readiness for return to sport (RTS) following anterior cruciate ligament (ACL) reconstruction is one of the most important responsibilities of physiotherapists involved in sports rehabilitation. Contemporary evidence indicates that RTS decisions should be based on objective functional criteria rather than time elapsed since surgery alone. Physiotherapy-based assessment focuses on evaluating muscle strength, limb symmetry, functional performance, balance, proprioception, neuromuscular control, and movement quality. Comprehensive evaluation of these factors helps reduce the risk of reinjury and promotes safe participation in sports.

**Muscle Strength Assessment:**


Restoration of lower-limb muscle strength is a fundamental requirement for successful return to sport. Following ACL reconstruction, significant weakness is commonly observed in the quadriceps and hamstring muscles due to pain, swelling, immobilization, and altered neuromuscular activation.

**Quadriceps Strength:**

Quadriceps strength is considered one of the most important predictors of functional recovery after ACL reconstruction. Persistent quadriceps weakness may impair dynamic knee stability, reduce athletic performance, and increase reinjury risk. Therefore, assessment of quadriceps strength is routinely incorporated into RTS testing protocols.

Isokinetic dynamometry is regarded as the gold standard for strength assessment. Hand-held dynamometry and functional strength tests may also be used when specialized equipment is unavailable.

Current evidence suggests that athletes should achieve at least 90% quadriceps strength symmetry compared with the uninvolved limb before returning to sport.




**Hamstring Strength:**

The hamstring muscles contribute significantly to dynamic stabilization of the knee joint by limiting anterior tibial translation. Adequate hamstring strength is particularly important following ACL reconstruction using hamstring tendon grafts, where postoperative weakness may persist.

Assessment of hamstring strength should be performed alongside quadriceps testing to ensure balanced muscle function and reduce injury risk.

**Limb Symmetry Index (LSI):**

The Limb Symmetry Index (LSI) is widely used to compare the performance of the reconstructed limb with the uninvolved limb.

**Calculation of LSI:**

LSI (%) = (Performance of Involved Limb ÷ Performance of Uninvolved Limb) × 100

An LSI score of 100% indicates complete symmetry between limbs. Most RTS guidelines recommend a minimum LSI of 90% before clearance for unrestricted sports participation.

The LSI can be applied to:

- Muscle strength measurements
- Hop test performance
- Balance assessments
- Functional movement tests

Although LSI is a valuable indicator of recovery, it should not be used in isolation because both limbs may experience deconditioning during rehabilitation.

**Functional Hop Tests:**
Functional hop tests are among the most commonly used assessments in RTS decision-making. These tests evaluate lower-limb power, stability, coordination, and confidence during dynamic activities.

**Single-Leg Hop Test:**
The single-leg hop test measures the maximum distance an athlete can hop forward on one leg while maintaining balance during landing.
Procedure:
- Athlete stands on one leg.
- Performs a maximal forward hop.
- Landing must be controlled and stable.
- Distance is measured from take-off to landing.

A minimum LSI of 90% is generally recommended for RTS clearance.

**Triple Hop Test:**
The triple hop test requires the athlete to perform three consecutive forward hops on the same limb.
This test assesses:
- Explosive strength
- Dynamic stability
- Neuromuscular control
- Lower-limb endurance

Triple hop performance provides additional information regarding an athlete's ability to repeatedly generate force while maintaining control.

**Crossover Hop Test:**
In the crossover hop test, the athlete performs three consecutive hops while crossing over a center line.
The test challenges:
- Dynamic balance
- Coordination
- Rotational control
- Multidirectional movement capability

Because many sports involve rapid directional changes, this assessment is particularly relevant for RTS evaluation.

**Six-Meter Timed Hop Test:**
The six-meter timed hop test measures the time required to hop a distance of six meters on one leg.
This test evaluates:
- Functional speed
- Agility
- Dynamic stability
- Lower-extremity power

Shorter completion times indicate better functional performance.

**Balance and Proprioception Assessment:**

Balance and proprioception are essential components of knee joint function. ACL injury disrupts mechanoreceptors within the ligament, resulting in impaired joint position sense and neuromuscular control.

Physiotherapists frequently assess balance using:

- Single-leg stance tests
- Star Excursion Balance Test (SEBT)
- Y-Balance Test
- Dynamic postural stability assessments

Athletes should demonstrate symmetrical balance performance and adequate postural control before RTS.

Improved proprioception contributes to:

- Enhanced movement efficiency
- Better joint stability
- Reduced injury risk
- Improved athletic performance

**Neuromuscular Control Assessment:**

Neuromuscular control refers to the coordinated activation of muscles responsible for maintaining joint stability during movement.

Deficits in neuromuscular control may lead to:

- Dynamic knee valgus
- Poor landing mechanics
- Abnormal movement patterns
- Increased ligament loading

Assessment methods include:

- Single-leg squat evaluation
- Step-down tests
- Jump-landing analysis
- Agility drills

Identification and correction of movement abnormalities are critical components of RTS preparation.

**Movement Quality Assessment:**

Movement quality assessment has gained increasing importance in sports rehabilitation. Athletes may achieve acceptable strength and hop test scores while continuing to exhibit poor movement patterns that increase reinjury risk.

Physiotherapists evaluate movement quality during:

**Jump Landing Tasks:**

Observations include:

- Knee alignment
- Hip control
- Trunk positioning
- Shock absorption strategies

**Single-Leg Squat Assessment:**

The single-leg squat provides valuable information regarding:

- Lower-limb alignment
- Dynamic stability
- Hip muscle function
- Neuromuscular control

Excessive knee valgus or trunk compensation may indicate inadequate readiness for RTS.

**Agility and Sport-Specific Testing:**

As rehabilitation progresses, athletes should undergo sport-specific functional assessments.

Examples include:

- Sprint drills
- Cutting maneuvers
- Direction-change tasks
- Reactive agility tests
- Acceleration and deceleration drills

These assessments simulate the physical demands encountered during competition and help determine sport-specific readiness.

**Recommended RTS Benchmarks:**

Based on current evidence, physiotherapists commonly consider the following criteria before granting RTS clearance:

| Assessment | Recommended Criterion |
|---|---|
| Knee Range of Motion | Full ROM |
| Quadriceps Strength | ≥90% LSI |
| Hamstring Strength | ≥90% LSI |
| Single-Leg Hop Test | ≥90% LSI |
| Triple Hop Test | ≥90% LSI |
| Crossover Hop Test | ≥90% LSI |
| Timed Hop Test | ≥90% LSI |
| Balance Assessment | Symmetrical Performance |
| Dynamic Stability | No Significant Deficits |
| Movement Quality | Normal Mechanics |

Athletes participating in high-level competitive sports may require even higher performance thresholds before RTS clearance.

**Clinical Implications for Physiotherapists:**

Physiotherapists play a central role in RTS decision-making by conducting comprehensive functional assessments and identifying residual impairments. Objective testing provides valuable information regarding recovery status and helps guide rehabilitation progression.

A criterion-based approach enables individualized treatment planning and minimizes the likelihood of premature return to sport. Integration of strength testing, functional performance evaluation, balance assessment, and movement quality analysis provides a comprehensive framework for RTS decision-making.

**Summary:**

Return-to-sport decisions following ACL reconstruction should be based on objective physiotherapy assessments rather than postoperative timelines alone. Muscle strength recovery, limb symmetry, hop test performance, balance, proprioception, neuromuscular control, and movement quality are key determinants of readiness for sports participation. Adoption of evidence-based RTS criteria can enhance rehabilitation outcomes, reduce reinjury risk, and support safe return to athletic performance.

**Current Evidence on Return-to-Sport Timing Following ACL Reconstruction:**

Determining the optimal timing for return to sport (RTS) following anterior cruciate ligament (ACL) reconstruction remains one of the most debated topics in sports rehabilitation. While advances in surgical techniques and physiotherapy rehabilitation have improved functional outcomes, the question of when an athlete can safely return to sport continues to challenge clinicians. Historically, RTS decisions were primarily based on postoperative timelines. However, recent evidence suggests that time alone is not an adequate indicator of recovery, and athletes should meet objective functional criteria before returning to competitive activities.

Current literature emphasizes that biological graft healing, restoration of muscle strength, neuromuscular control, functional performance, and movement quality require sufficient time to recover. Consequently, RTS timing should be viewed as one component of a broader criterion-based decision-making process.


**Biological Healing Following ACL Reconstruction:**

Successful return to sport depends partly on the biological incorporation and maturation of the graft. Following reconstruction, the graft undergoes a process known as ligamentization, during which the transplanted tissue gradually acquires characteristics similar to a native ligament.

This process involves:

- Initial graft necrosis
- Cellular repopulation
- Revascularization
- Collagen remodeling
- Structural maturation

Although functional improvements may occur within a few months after surgery, biological healing continues for many months. Research suggests that complete graft maturation may require 12 months or longer, supporting a cautious approach to RTS decision-making.




**Return to Sport Before 9 Months:**

Many athletes attempt to return to sports within the first 6 to 9 months after surgery, particularly in competitive environments where pressure to resume participation is high. However, evidence consistently demonstrates that early RTS is associated with an increased risk of reinjury.

Several studies have reported that athletes returning before 9 months often exhibit:

- Persistent quadriceps weakness
- Reduced hamstring strength
- Incomplete neuromuscular recovery
- Deficits in hop performance
- Abnormal movement mechanics

These impairments may compromise dynamic knee stability and increase stress on the reconstructed ligament.

Furthermore, younger athletes participating in pivoting and cutting sports appear particularly vulnerable to reinjury when returning prematurely. The combination of high athletic demands and incomplete recovery creates a substantial risk for graft rupture and secondary injuries.

Therefore, most contemporary rehabilitation guidelines discourage unrestricted RTS before 9 months unless objective functional criteria have been successfully achieved.

**Return to Sport Between 9 and 12 Months:**

The period between 9 and 12 months following ACL reconstruction is commonly considered the most appropriate timeframe for RTS assessment. During this stage, athletes generally demonstrate substantial improvements in:

- Quadriceps strength
- Hamstring strength

- Functional hop performance
- Dynamic balance
- Sport-specific skills

Many rehabilitation programs use this period to conduct comprehensive RTS testing and evaluate readiness for competitive participation.

Research indicates that delaying RTS until at least 9 months after surgery may significantly reduce reinjury risk. Athletes who achieve acceptable strength symmetry, functional performance, and movement quality during this period generally demonstrate better long-term outcomes than those who return earlier.

Nevertheless, not all athletes recover at the same rate. Some individuals may continue to exhibit significant functional deficits despite reaching the 9-month milestone. Consequently, physiotherapists should prioritize objective assessment findings over postoperative timelines when making RTS recommendations.

**Return to Sport After 12 Months:**

Recent evidence increasingly supports delaying RTS until 12 months or longer in athletes participating in high-risk sports. This approach allows additional time for:

- Graft maturation
- Strength development
- Neuromuscular adaptation
- Movement retraining
- Sport-specific conditioning

Athletes returning after 12 months often demonstrate superior functional outcomes and lower rates of secondary ACL injury. Extended rehabilitation also provides opportunities to address residual deficits that may not be detected during routine clinical assessment.

High-level athletes involved in sports requiring frequent cutting, pivoting, and jumping may particularly benefit from a longer rehabilitation period before RTS clearance.

However, prolonged delays should not be based solely on time. Athletes who have successfully achieved all RTS criteria and demonstrate excellent functional performance may safely return before 12 months under appropriate clinical supervision.

**Time-Based Versus Criterion-Based Approaches:**

Traditional RTS models relied heavily on fixed timelines, assuming that all athletes recover at a similar rate. This approach fails to account for individual differences in healing, rehabilitation progress, and functional recovery.

In contrast, criterion-based rehabilitation emphasizes achievement of specific performance benchmarks before advancement. This model evaluates:

- Muscle strength recovery
- Limb symmetry
- Functional performance
- Dynamic stability
- Movement quality
- Sport-specific readiness

Current evidence strongly supports criterion-based decision-making as a more reliable strategy for reducing reinjury risk and optimizing athletic performance.

**Factors Influencing RTS Timing:**

Several factors influence the timing of return to sport following ACL reconstruction:

**Age:**

Younger athletes often demonstrate greater physical recovery potential but may also exhibit higher reinjury rates due to increased participation in high-demand sports.

**Type of Sport:**

Athletes involved in cutting and pivoting sports typically require more comprehensive rehabilitation compared with participants in straight-line activities such as running or cycling.

**Graft Type:**


Different graft choices may influence rehabilitation progression and muscle strength recovery.

**Functional Recovery:**

Achievement of strength and functional performance benchmarks remains one of the most important determinants of RTS timing.

**Rehabilitation Compliance:**

Athletes who adhere consistently to rehabilitation programs generally achieve better outcomes and may progress more efficiently through RTS stages.




**Physiotherapy Recommendations:**

Based on current evidence, physiotherapists should avoid making RTS decisions solely on postoperative timelines. Instead, RTS clearance should be based on a combination of:

- Minimum postoperative healing time

- Objective strength testing
- Functional hop test performance
- Balance assessment
- Movement quality evaluation
- Sport-specific testing

A multidisciplinary approach involving physiotherapists, orthopedic surgeons, strength and conditioning specialists, and sports medicine professionals can further enhance decision-making and athlete safety.

**Summary:**

The timing of return to sport following ACL reconstruction remains a critical factor influencing rehabilitation outcomes and reinjury risk. Although many athletes return to sports between 9 and 12 months after surgery, current evidence supports a criterion-based approach rather than reliance on time alone. Athletes who demonstrate adequate strength, functional performance, balance, neuromuscular control, and movement quality are more likely to achieve successful and safe return to sport. Delaying RTS until objective recovery criteria are met may reduce reinjury risk and improve long-term athletic performance.

**References**

1. Almeida, V. F., Silva, R. M., & Santos, L. P. (2026). Criteria for return to sport after anterior cruciate ligament reconstruction: A literature review. Research, Society and Development, 15(2), e10215251050.
2. Ardern, C. L., Glasgow, P., Schneiders, A., Witvrouw, E., Clarsen, B., Cools, A., Gojanovic, B., Griffin, S., Khan, K. M., Moksnes, H., Phillips, N., Reurink, G., Sadler, R., Silbernagel, K. G., Thorborg, K., Wangensteen, A., Wilk, K. E., & Bizzini, M. (2020). 2016 Consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. British Journal of Sports Medicine, 54(14), 853–864.
3. Brinlee, A. W., Dickenson, S. B., Hunter-Giordano, A., & Snyder-Mackler, L. (2022). ACL reconstruction rehabilitation: Clinical data, biologic healing, and criterion-based milestones to inform a return-to-sport guideline. Sports Health, 14(3), 365–376.
4. Davies, G. J., McCarty, E., Provencher, M., & Manske, R. C. (2020). ACL return-to-sport guidelines and criteria. Current Reviews in Musculoskeletal Medicine, 13(3), 307–314.
5. Drole, K., & Paravlic, A. H. (2022). Interventions for increasing return-to-sport rates after anterior cruciate ligament reconstruction surgery: A systematic review. Frontiers in Psychology, 13, 940092.
6. Foley, A., Richter, C., & Moran, R. (2025). Return to sport following ACL reconstruction: Current concepts and clinical considerations. Current Reviews in Musculoskeletal Medicine, 18(4), 215–228.
7. Glattke, K. E., Tummala, S. V., & Chhabra, A. (2022). Anterior cruciate ligament reconstruction recovery and rehabilitation: A systematic review. The Journal of Bone and Joint Surgery, 104(8), 739–752.
8. Gokeler, A., Dingenen, B., & Hewett, T. E. (2022). Rehabilitation and return-to-sport testing after anterior cruciate ligament reconstruction: Where are we in 2022? Arthroscopy, Sports Medicine, and Rehabilitation, 4(1), e77–e82.
9. Golberg, E., Hammoud, S., & Capin, J. J. (2024). Anterior cruciate ligament reconstruction return-to-sport assessment: A scoping review. Sports Health, 16(2), 145–156.

10. Grindem, H., Arundale, A. J. H., & Ardern, C. L. (2020). Evidence-based recommendations for return to sport after ACL reconstruction. British Journal of Sports Medicine, 54(20), 1223–1230.
11. Joreitz, R., Nagelli, C., & Hewett, T. E. (2025). Sport-specific training and return to sport after ACL reconstruction in elite athletes: A narrative review. Annals of Joint, 10(4), 1–14.
12. Katasani, S. B. R. (2026). An integrated biopsychosocial, nutritional and administrative approach to rehabilitation outcomes: Implications for physiotherapy practice. International Journal of Pharmaceutical Sciences, 4(2), 112–120.
13. Katasani, S. B. R., Kumar, A., Stanly, W., & Shekhar, P. C. (2026). Handbook of Cupping Therapy: Physiotherapy and Occupational Therapy. Academic Publishers.
14. Kotsifaki, R., Korakakis, V., King, E., Barbosa, O., Maree, D., Pantouveris, M., Bjerregaard, A., Luomajoki, J., Wilhelmsen, J., & Whiteley, R. (2023). Aspetar clinical practice guideline on rehabilitation after anterior cruciate ligament reconstruction. British Journal of Sports Medicine, 57(9), 500–514.
15. Mardani-Kivi, M., Azari, Z., & Hasannejad, F. (2020). Return to sport activity after anterior cruciate ligament reconstruction: A 6–10 years follow-up study. Journal of Clinical Orthopaedics and Trauma, 11(Suppl. 3), S319–S324.
16. Thomeé, R., Kaplan, Y., Kvist, J., Myklebust, G., Risberg, M. A., Theisen, D., Toksvig-Larsen, S., & Witvrouw, E. (2020). Muscle strength and hop performance criteria prior to return to sports after ACL reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 28(5), 1798–1805.
17. Waldron, K., Brown, M., Calderon, A., & Feldman, M. (2022). Anterior cruciate ligament rehabilitation and return to sport: How fast is too fast? Arthroscopy, Sports Medicine, and Rehabilitation, 4(1), e175–e179.
18. Welling, W., Benjaminse, A., Lemmink, K., & Gokeler, A. (2020). Passing return-to-sports tests after ACL reconstruction is associated with greater likelihood for return to sport but fails to identify second injury risk. The Knee, 27(3), 949–957.
19. Xu, S., Smith, T., Johnson, M., & Brown, K. (2023). Return to sport after primary anterior cruciate ligament reconstruction: A survey of current clinical practice. Kansas Journal of Medicine, 16(1), 105–112.