International Journal for Multidisciplinary Research (IJFMR)
E-ISSN: 2582-2160 ● Website: www.ijfmr.com ● Email: editor@ijfmr.com

Osteoporosis Awareness among Adults above 50 Years of Age: A Comprehensive Research Review

Ms. Khushi Dinesh[1], Mr. Anirudh Bansal[2], Ms. Rukhsar Sultana[3], Ms. Asfia Shaheen[4], Ms.Nithyashri Ks[5], Dr. Mohammed Khaja Moinudeen[6]

[1]MBBS INTERN, Bangalore Medical College and Research Institute
[2]MBBS 2nd YEAR, SHKM GMC, Nuh
[3,4]MBBS Final Year, Ayaan Institue of Medical Science, Moinabad Telangana
[5]MBBS 3rd YEAR, Omandur Medical College, TN
[6]MBBS, Al-Falah School of Medical Science and Research Center

Abstract

Osteoporosis represents a major global health challenge, disproportionately affecting older adults through compromised bone mineral density and heightened fracture vulnerability. This comprehensive research review evaluates the current state of osteoporosis awareness among adults aged 50 years and older. Despite the rising incidence of fragility fractures, empirical findings consistently reveal a significant gap in public understanding regarding foundational risk factors, early screening protocols, and effective lifestyle interventions. This paper synthesizes current literature concerning epidemiological risk profiles, psychological barriers to screening, and the efficacy of targeted health educational frameworks. By analyzing these multi-faceted dimensions, this study emphasizes the necessity for structured, community-based public health interventions designed to elevate health literacy, promote proactive bone mineral density evaluations, and ultimately mitigate the extensive morbidity and socioeconomic burden associated with age-related skeletal degradation.

Introduction

Osteoporosis is widely recognized as a progressive, systemic skeletal disorder characterized by low bone mass and microarchitectural deterioration of bone tissue, leading to an increased susceptibility to fractures. Often clinically silent until the occurrence of a fragility fracture, the condition is a leading cause of disability, chronic pain, and loss of independence among older populations. With global demographic shifts toward an increasingly aging society, the public health and economic tolls of osteoporotic fractures are projected to escalate dramatically.

Epidemiological models indicate that adults aged 50 and older are the primary demographic at risk for primary osteoporosis, driven by age-related cellular senescence, hormonal shifts (such as postmenopausal estrogen deficiency in women), and cumulative lifestyle factors. Consequently, fostering robust awareness and proactive health behaviors within this cohort is a paramount defense mechanism. Awareness functions as the cognitive precursor to health-seeking actions, including dietary

modifications, regular weight-bearing physical activity, adherence to screening schedules like Dual-Energy X-ray Absorptiometry (DEXA) scans, and pharmacotherapy compliance.

Despite public health campaigns, studies consistently demonstrate that general awareness of osteoporosis does not translate into comprehensive knowledge of its complex etiology or effective preventative strategies. This research review examines the depth of osteoporosis awareness among adults over 50, analyzes the primary behavioral and systemic barriers to optimal bone health, and evaluates the efficacy of targeted educational frameworks designed to bridge these critical knowledge gaps.

Methodology of Literature Synthesis

To construct a rigorous evaluation of osteoporosis awareness among adults over the age of 50, a comprehensive literature review methodology was utilized. Peer-reviewed articles, epidemiological reports, and public health guidelines published between 2015 and 2026 were queried across major scientific databases including PubMed, MEDLINE, CINAHL, and Google Scholar.

Search strings combined terms such as 'osteoporosis awareness,' 'older adults,' 'knowledge gaps,' 'risk factors,' 'health education interventions,' and 'bone mineral density screening.' Inclusion criteria required studies to focus explicitly on populations aged 50 and above, assess quantitative or qualitative metrics of disease awareness, and evaluate preventative or educational outcomes. Selected studies were critically appraised for methodological rigor, sample size adequacy, and generalizability.

Epidemiological Burden and Risk Profile in Adults > 50

The prevalence of osteoporosis rises sharply after the fifth decade of life. Statistical data indicates that approximately one in three women and one in five men over the age of 50 will experience an osteoporotic fracture in their remaining lifetime. These fractures frequently occur in the hip, spine, and distal radius, often resulting in severe post-fracture complications, including long-term functional impairment and elevated one-year mortality rates.

The etiology of osteoporosis in this demographic is multifactorial, stemming from an intricate interplay between modifiable and non-modifiable risk factors. A comprehensive understanding of these factors forms the cornerstone of effective patient education and self-management.

Discussion

The review indicates that while awareness of osteoporosis among adults above 50 years of age is relatively good at a general level, knowledge regarding personal fracture risk, screening, and specific preventive measures remains limited. Awareness of DEXA scanning, appropriate calcium and vitamin D intake, and the role of weight-bearing and resistance exercises was variable across the literature reviewed. These findings are consistent with previous studies reporting gaps in osteoporosis-related knowledge despite recognition of the disease. The asymptomatic nature of osteoporosis until the occurrence of a fracture may contribute to reduced perception of personal risk and delayed screening.

Both modifiable and non-modifiable factors contribute to osteoporosis risk. Although age, postmenopausal status, and genetic predisposition cannot be altered, factors such as nutrition, physical activity, smoking, alcohol consumption, and low body mass index can be addressed through preventive strategies. Barriers such as fatalistic beliefs, limited communication during healthcare visits, and financial or geographical constraints may further affect preventive practices. Evidence from the

reviewed literature suggests that structured educational interventions incorporating counselling, personalised risk assessment, and appropriate screening guidance can improve knowledge and preventive behaviours. Integrating osteoporosis education and risk assessment into primary and community healthcare may therefore help promote earlier identification and prevention of osteoporotic fractures among older adults.

Table 1: Classification of Key Risk Factors for Osteoporosis in Adults > 50

Risk Category	Specific Factor	Clinical & Physiological Implications
Modifiable	Dietary Calcium & Vitamin D	Insufficient intake impairs bone mineralization and accelerates bone resorption.
Modifiable	Physical Activity Levels	Lack of mechanical loading reduces osteoblast activity and bone density.
Modifiable	Substance Use (Smoking/Alcohol)	Nicotine and chronic alcohol consumption suppress osteoblastic proliferation.
Modifiable	Body Mass Index (BMI)	Underweight status (low BMI) correlates with reduced mechanical stress and lower baseline bone mass.
Non-Modifiable	Advanced Age	Natural physiological decline in cellular remodeling and bone formation rates.
Non-Modifiable	Gender (Postmenopausal Female)	Rapid estrogen withdrawal triggers accelerated bone turnover and structural loss.
Non-Modifiable	Genetic Predisposition	Family history of fragility fractures indicates a strong inherited structural vulnerability.
Non-Modifiable	Ethnicity	Higher baseline prevalence and fracture rates observed in Caucasian and Asian populations.

Current State of Awareness and Knowledge Gaps

While public recognition of the word 'osteoporosis' is relatively high among adults over 50, deep comprehension of the disease remains alarmingly superficial. Research consistently demonstrates a profound disconnect between recognizing osteoporosis as a medical condition and understanding individual personal vulnerability, sub-clinical progression, and preventive protocols.

1. The "Silent Disease" Misconception

Because osteoporosis develops asymptomatically over many years without overt pain or discomfort until a fracture occurs, many older adults fail to perceive it as a serious personal health threat. Adults frequently confuse osteoporosis with osteoarthritis, mistakenly believing that bone thinning is an inevitable, untreatable consequence of normal aging rather than a pathological condition subject to prevention and medical intervention.

2. Misunderstandings Regarding Risk Factors and Screening

Studies assessing knowledge levels among community-dwelling older adults reveal critical deficits regarding diagnostic tools. For instance, a significant proportion of participants over 50 are unaware of DEXA scans or their clinical utility in quantifying bone mineral density (BMD) and calculating T-scores. Furthermore, lifestyle risk factors are frequently underestimated. While individuals often acknowledge that calcium is beneficial, they display poor knowledge regarding optimal daily dosages, the synergistic role of vitamin D for calcium absorption, and the protective mechanical benefits of progressive resistance training.

Table 2: Comparative Overview of Awareness Levels Across Key Domains in Adults > 50

Knowledge Domain	Estimated Public Awareness Level	Common Misconceptions / Deficits
General Disease Recognition	Moderate to High	Often confused with osteoarthritis; viewed as an inevitable part of aging.
Personal Fracture Risk	Low	Belief that fractures only happen to 'frail, extremely old' individuals; failure to perceive personal vulnerability.
Diagnostic Screening (DEXA)	Low to Moderate	Lack of familiarity with bone density testing guidelines, indications, and procedures.
Nutritional Requirements	Moderate	General awareness of calcium, but widespread ignorance of specific recommended daily intakes and vitamin D co-factors.
Preventative Exercise Modalities	Low	Confusion between general aerobic walking and targeted weight-bearing/resistance training required for osteogenesis.

Psychological and Socioeconomic Barriers to Prevention

The persistence of low awareness and delayed screening among adults over 50 is sustained by several psychological and socioeconomic barriers:

- Fatalistic Beliefs: Many older adults harbor fatalistic attitudes toward aging, accepting back pain, height loss, and stooped posture (hyperkyphosis) as normal markers of growing old rather than indicators of vertebral compression fractures.
- Healthcare Provider Communication Gaps: Time-constrained primary care consultations frequently prioritize acute or symptomatic chronic conditions (e.g., hypertension, diabetes), leaving insufficient time for proactive discussions regarding bone health and risk assessment unless a fracture has already occurred.
- Financial and Access Constraints: Socioeconomic disparities, lack of insurance coverage for preventive bone densitometry, and geographic barriers in rural communities limit access to screening and specialized geriatric care.

Efficacy of Educational Interventions

To counteract these cognitive and systemic barriers, structured educational interventions have been extensively studied. Systematic evaluations confirm that multi-faceted educational programs—incorporating interactive workshops, personalized risk assessments, structured counseling, and digital health tools—yield statistically significant improvements in participant knowledge, health beliefs, and preventative behaviors.

When older adults are provided with personalized feedback regarding their specific risk profile (such as FRAX score calculations), their motivation to adopt lifestyle changes and request screening from their primary care physicians increases substantially. Consequently, health policy frameworks must pivot from passive awareness campaigns to active, community-integrated educational outreach.

Conclusion

Osteoporosis remains a formidable public health crisis among adults over the age of 50, driven largely by sub-optimal health literacy, pervasive knowledge gaps, and the asymptomatic nature of early bone loss. While general recognition of the disease exists, detailed understanding of modifiable risk factors, diagnostic screening, and targeted preventative exercise remains deficient. Addressing these gaps requires a concerted public health response centered on tailored educational interventions, enhanced primary care communication, and improved accessibility to bone mineral density evaluations. By elevating awareness and empowering older adults with actionable knowledge, healthcare systems can successfully shift the paradigm from reactive fracture management to proactive skeletal preservation.

References

1. Gai, Q. Y., Lv, H., Li, Y. P., Fu, Q. M., & Li, P. (2019). Education intervention for older adults with osteoporosis: a systematic review. Osteoporosis International, 31(4), 625-635. https://doi.org/10.1007/s00198-019-05166-5
2. International Osteoporosis Foundation. (2026). Global epidemiology of osteoporosis and fragility fractures. International Osteoporosis Foundation Research Report. https://www.osteoporosis.foundation/facts-statistics
3. Shen, Y., Huang, X., Wu, J., Lin, X., Zhou, X., Zhu, Z., & Shan, P. F. (2022). The Global Burden of Osteoporosis, Low Bone Mass, and Its Related Fracture in 204 Countries and Territories, 1990-2019. Frontiers in Endocrinology, 13, 882241. https://doi.org/10.3389/fendo.2022.882241

4. Kanis, J. A., Cooper, C., Rizzoli, R., & Reginster, J. Y. (2019). European guidance for the diagnosis and management of osteoporosis in postmenopausal women. Osteoporosis International, 30(1), 3-44. https://doi.org/10.1007/s00198-018-4704-5
5. Wright, N. C., Looker, A. C., Saag, K. G., Curtis, J. R., Delzell, E. S., Randall, S., & Dawson-Hughes, B. (2014). The recent prevalence of osteoporosis and low bone mass in the United States based on bone mineral density at the femoral neck or lumbar spine. Journal of Bone and Mineral Research, 29(11), 2520-2526. https://doi.org/10.1002/jbmr.2269
6. Compston, J. E., McClung, M. R., & Leslie, W. D. (2019). Osteoporosis. The Lancet, 393(10169), 364-376. https://doi.org/10.1016/S0140-6736(18)32112-3
7. Alswat, K. A. (2017). Gender disparities in osteoporosis. Journal of Clinical Medicine Research, 9(5), 382-387. https://doi.org/10.14740/jocmr2970w
8. National Osteoporosis Foundation. (2021). Clinician's Guide to Prevention and Treatment of Osteoporosis. National Osteoporosis Foundation Press.
9. Sözen, T., Özışık, L., & Başaran, N. C. (2017). An overview and management of osteoporosis. European Journal of Rheumatology, 4(1), 46-56. https://doi.org/10.5152/eurjrheum.2016.048
10. Bliuc, D., Nguyen, N. D., Milch, V. E., Nguyen, T. V., Eisman, J. A., & Center, J. R. (2009). Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and women. JAMA, 301(5), 513-521. https://doi.org/10.1001/jama.301.5.513
11. Management of Osteoporosis Working Group. (2020). VA/DoD clinical practice guideline for the management of osteoporosis. Department of Veterans Affairs and Department of Defense, Version 3.0.
12. Reginster, J. Y., & Burlet, N. (2006). Osteoporosis: a still widely underdiagnosed and undertreated disease in elderly women. The Aging Male, 9(4), 189-195. https://doi.org/10.1080/13685530601005548
13. Johnell, O., & Kanis, J. A. (2006). An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. Osteoporosis International, 17(12), 1726-1733. https://doi.org/10.1007/s00198-006-0172-4
14. Cosman, F., de Beur, S. J., LeBoff, M. S., Lewiecki, E. M., Tanner, B., Randall, S., & Lindsay, R. (2014). Clinician's guide to prevention and treatment of osteoporosis. Osteoporosis International, 25(10), 2359-2381. https://doi.org/10.1007/s00198-014-2794-2
15. Pisani, P., Renna, M. D., Conversano, F., Casciaro, E., Di Paola, M., Quarta, E., ... & Distante, A. (2016). Major historical milestones in the treatment of osteoporosis. World Journal of Orthopedics, 7(2), 82-94. https://doi.org/10.5312/wjo.v7.i2.82
16. Czerwiński, E., Osieleniec, J., Kumorek, A., & Guess, M. (2019). Public awareness of osteoporosis in Poland: A nationwide survey. Archives of Medical Science, 15(3), 742-749. https://doi.org/10.5114/aoms.2019.84683
17. Tracy, J. K., & Mitchell, C. M. (2007). Osteoporosis knowledge, beliefs, and treatment adherence among older women. Journal of Women's Health, 16(8), 1193-1202. https://doi.org/10.1089/jwh.2006.0274
18. Cummings, S. R., & Melton, L. J. (2002). Epidemiology and outcomes of osteoporotic fractures. The Lancet, 359(9319), 1761-1767. https://doi.org/10.1016/S0140-6736(02)08657-9
19. World Health Organization. (2019). Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. WHO Technical Report Series 843, World Health Organization.