

Language Issues in Autistic Children: Characteristics, Assessment and Intervention: A Brief Narrative Review

Lavkesh

Assistant Professor, Sarthi Institute of Developmental Studies, Barabanki, Uttar Pradesh

Abstract

Language and communication differences are among the most clinically significant features observed in autistic children. However, the language profile of autism is highly heterogeneous. Some children develop fluent spoken language with age-appropriate vocabulary and grammar, others show significant receptive and expressive language difficulties, and some remain minimally speaking and communicate more effectively through gestures or augmentative and alternative communication (AAC). Difficulties may involve pragmatics, vocabulary and semantics, grammar and syntax, narrative skills, prosody, receptive language and the flexible use of language in social situations. Echolalia, repetitive language, unusual prosody and highly focused vocabulary may also occur. This brief narrative review examines major language issues associated with autism, their functional consequences, assessment principles, and evidence-informed intervention. Current evidence supports individualized, functional and family-centred approaches that consider both spoken and nonspoken communication. Early assessment by speech-language professionals, evaluation of hearing, attention to social communication, and provision of AAC when indicated may improve participation and communication outcomes.

Keywords: Autism spectrum disorder; autistic children; language disorder; pragmatic language; echolalia; social communication; AAC; speech-language therapy.

1. Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental condition associated with differences in social communication and patterns of restricted or repetitive behaviour and interests. Communication development among autistic children varies substantially. A child may have little functional speech, another may use sentences but struggle to understand spoken instructions, while another may possess an extensive vocabulary yet experience difficulty maintaining reciprocal conversation or interpreting another person's intended meaning. NIDCD describes differences in language development, repetitive or rigid language, conversational interaction and nonverbal communication among autistic children. Contemporary research therefore discourages viewing autism as producing a single language profile. A review of language in autism identified at least three broad patterns: autistic individuals without major structural-language impairment, those with accompanying structural-language impairment, and minimally verbal individuals. A 2026 systematic review similarly found considerable heterogeneity in phonology, morphology, syntax and semantics among autistic children, particularly when autism occurs with a co-existing language impairment.

2. Major Language Issues in Autistic Children

2.1 Receptive and Expressive Language

Receptive language refers to understanding words, sentences, questions and instructions, whereas expressive language refers to communicating ideas through words, sentences or other modes. In autistic children, these abilities may develop unevenly. Some children can label many objects or recite familiar material but have difficulty understanding abstract questions, complex instructions or language presented outside a familiar context.

Expressive difficulties may include delayed first words, restricted vocabulary, short utterances, grammatical difficulties or limited spontaneous language. Importantly, the apparent amount of speech should not automatically be considered equivalent to the child's level of language comprehension.

2.2 Pragmatic and Social-Communication Difficulties

Pragmatics concerns how language is used socially and appropriately in context. It is frequently an important area of difficulty in autism. Children may experience difficulty initiating interaction, taking conversational turns, maintaining a shared topic, modifying information according to the listener, interpreting indirect requests or understanding humour, sarcasm and figurative expressions. Systematic-review evidence indicates that pragmatic-language difficulties are particularly prominent in autism, although substantial individual variation exists.

ASHA recommends considering joint attention, communicative initiation, social reciprocity, gestures, facial expression, prosody, topic management and conversational turn-taking when evaluating social communication.

2.3 Echolalia and Repetitive Language

Some autistic children repeat words, phrases or longer utterances previously heard. **Immediate echolalia** occurs soon after hearing an utterance, whereas **delayed echolalia** may reproduce language heard much earlier from family members, media or familiar situations. Repetitive expressions and scripts should not automatically be considered meaningless. Their communicative function should be examined because a repeated phrase may serve purposes such as requesting, protesting, regulating emotion, participating in interaction or recalling a familiar routine.

2.4 Vocabulary, Grammar and Narrative Skills

Structural language is also variable. Some autistic children demonstrate relatively strong vocabulary while experiencing difficulties with grammatical morphology, complex sentence structures or comprehension. The latest systematic evidence indicates overlap between autistic children with language impairment and children with developmental language disorder in areas such as receptive and expressive language, morphosyntax and complex sentence processing.

Narrative language can also be challenging. A child may identify facts or events but have difficulty organising them into a coherent story, identifying the listener's informational needs, explaining cause-and-effect relationships or describing characters' intentions and emotions.

Table 1. Common Language Issues and Possible Functional Effects

Language domain	Possible characteristics	Functional effect
Receptive language	Difficulty understanding questions, instructions or abstract language	Problems following classroom and daily routines
Expressive language	Limited spontaneous vocabulary, short or grammatically immature sentences	Difficulty explaining needs, experiences and ideas

Pragmatics	Difficulty with turn-taking, topic maintenance and implied meaning	Reduced effectiveness of social conversation
Echolalia/scripts	Repetition of previously heard language	May function as communication but be misunderstood by others
Semantics	Literal interpretation or difficulty with flexible word meanings	Difficulty with figurative/ambiguous language
Narrative	Reduced coherence or organisation of events	Academic and social communication difficulties
Prosody	Differences in rhythm, stress, pitch or intonation	Listener may misinterpret communicative intent
Nonverbal communication	Differences in gesture, joint attention or facial expression	Spoken communication may be harder for partners to interpret

3. Assessment

Language assessment in autism should be comprehensive rather than based only on the number of words spoken. The speech-language pathologist should examine receptive and expressive language, semantics, morphology and syntax, narrative ability, pragmatic communication, speech sound production, prosody and functional communication across different environments.

Naturalistic observation and language sampling are particularly valuable because standardized tests alone may not demonstrate how effectively a child communicates at home, school or during interaction with peers. Information from parents and teachers should therefore complement formal testing. Hearing evaluation is also important because reduced responses to spoken communication can occur in both hearing difficulties and autism, and the two may coexist.

For multilingual children, autism should not be attributed simply to exposure to more than one language. Assessment should consider the languages actually used within the child's family, school and community.

4. Language Intervention

Intervention should aim primarily to improve **functional communication and participation**, rather than simply increasing the number of spoken words. Goals may include requesting, commenting, refusing appropriately, sharing information, answering and asking questions, understanding instructions, expanding vocabulary and grammar, developing narratives and improving conversational reciprocity.

Naturalistic language teaching, shared activities, visual supports, explicit vocabulary instruction and intervention within meaningful routines may be used according to the child's profile. ASHA notes that language intervention for autistic individuals may range from structured clinician-directed approaches to naturalistic developmental approaches and should reflect individual communication needs and strengths. Children who have little or unreliable functional speech should also have access to **augmentative and alternative communication (AAC)** when appropriate. AAC may include gestures, communication boards, picture-based systems, tablets or speech-generating devices. Evidence from a systematic review and meta-analysis indicates that naturalistic developmental behavioural interventions and aided AAC can support language development in autistic children with minimal speech. Intervention is generally most useful when parents, teachers and other regular communication partners are involved so that skills are practised across everyday settings.

5. Conclusion

Language difficulties in autistic children cannot be described by a single pattern. They may involve receptive or expressive language, pragmatics, semantics, grammar, narrative organisation, prosody, echolalia or nonverbal communication, with considerable variation between children. A child with sophisticated vocabulary may still experience substantial functional communication difficulties, while a minimally speaking child may have meaningful communicative abilities that are underestimated when assessment focuses exclusively on speech. Effective management therefore requires individualized and multidisciplinary assessment, including hearing evaluation and detailed speech-language assessment. Intervention should focus on meaningful communication, participation and the child's individual strengths and needs. Appropriate speech-language therapy, communication-partner training, family involvement and AAC can provide autistic children with more effective ways to express choices, ideas, emotions and needs.

References

1. National Institute on Deafness and Other Communication Disorders. *Autism Spectrum Disorder: Communication Problems in Children*. National Institutes of Health.
2. American Speech-Language-Hearing Association. *Autism and Autism Spectrum Disorder: Practice Portal*. ASHA.
3. American Speech-Language-Hearing Association. *Spoken Language Disorders: Practice Portal*. ASHA.
4. Doghadze I, Gagoshidze T. Structural language profiles in children with autism spectrum disorder and developmental language disorder: a systematic review. *Frontiers in Psychology*. 2026;17:1920573. doi:10.3389/fpsyg.2026.1920573.
5. Vogindroukas I, et al. Developmental Language Disorder and Autism Spectrum Disorder: similarities in pragmatic language abilities—a systematic review. *International Journal of Developmental Disabilities*.
6. Georgiou N, Spanoudis G. Language in autism: domains, profiles and co-occurring conditions. *Journal of Neural Transmission*. 2023. doi:10.1007/s00702-023-02592-y.
7. Pope L, Light J, Laubscher E. The effect of naturalistic developmental behavioral interventions and aided AAC on the language development of children on the autism spectrum with minimal speech: a systematic review and meta-analysis. *Journal of Autism and Developmental Disorders*. 2024. doi:10.1007/s10803-024-06382-7.