

Postpartum Depression: A Narrative Critical Literature Review of Homeopathic Therapeutics with Special Reference to Sepia, Natrum Muriaticum, and Pulsatilla

Dr. Lekha Rajesh¹, Dr. Hemanth Babu R²

¹Doctor, MSc Psychology Student, Centre for Distance and Online Education (CDOE)
JAIN (Deemed-to-be University), Bengaluru, India

²Professorr Department of Psychology, Centre for Distance and Online Education (CDOE)
JAIN (Deemed-to-be University), Bengaluru, India

Abstract

Postpartum Depression (PPD) represents a profound maternal mental health challenge affecting emotional well-being, mother-infant bonding, and family stability worldwide. While conventional psychiatric and psychotherapeutic interventions serve as the primary standard of care, complementary modalities such as homeopathy are frequently sought by mothers navigating perinatal vulnerabilities due to concerns surrounding medication safety during lactation and personal treatment preferences. This paper presents a narrative critical literature review examining the theoretical descriptions and clinical relevance of three classical homeopathic remedies-Sepia officinalis, Natrum muriaticum, and Pulsatilla nigricans- in addressing postpartum emotional disturbances.

From the daily clinical perspective of practicing homeopathic physicians, individualized prescribing based on symptom similarity, constitutional typology, and mental generals consistently yields positive therapeutic outcomes in patient recovery. However, a significant gap persists in the medical literature: homeopathic practitioners are frequently unable to produce large-scale clinical trials or robust empirical research papers due to systemic challenges including lack of standardized documentation, limited research infrastructure, and patient participation barriers. This article critically examines this limitation, advocates for rigorous clinical documentation, and calls upon the medical and scientific community to bridge traditional clinical success with modern evidence-based research methodologies, ensuring that the benefits of homeopathic care are scientifically validated and accessible to a broader community.

Keywords: Postpartum Depression, Homeopathy, Sepia, Natrum Muriaticum, Pulsatilla, Individualized Prescribing, Clinical Documentation, Evidence-Based Research,

1. Introduction

1.1 Background of the Study & Clinical Context

Postpartum Depression (PPD) is a major public health concern transcending cultural, socioeconomic, and geographical boundaries. Defined as a depressive episode occurring within the first year after childbirth, PPD manifests through persistent sadness, irritability, severe fatigue, impaired maternal-infant bonding,

and marked disturbances in daily functioning (O'Hara & McCabe, 2013). Globally, prevalence rates range between 10-20%, with higher incidence reported in low- and middle-income countries due to socioeconomic stressors, limited healthcare access, and cultural stigma (WHO, 2020).

In India, maternal mental health remains under-recognized, with PPD often dismissed as normal adjustment' or overshadowed by cultural expectations surrounding motherhood (Patel et al., 2012). Conventional management relies on pharmacological interventions (e.g., SSRIs) and psychotherapies (CBT, IPT). However, barriers such as societal stigma, reluctance to take psychotropic medications during breastfeeding, and limited healthcare access drive patients toward complementary systems.

In daily clinical practice, homeopathic physicians frequently encounter and treat a substantial number of PPD patients. By applying the fundamental principle of individualization-matching the patient's unique totality of physical, mental, and constitutional symptoms to the most similar remedy (the similimum) clinicians observe meaningful clinical recoveries and emotional stabilization. Yet, despite these positive clinical observations, the homeopathic community faces a critical limitation: the scarcity of large-scale, published evidence-based research papers.

1.2 Statement of the Research Problem & Rationale

The core paradox of homeopathic practice in modern healthcare lies in the divergence between positive daily clinical outcomes and the paucity of formal, large-scale empirical documentation. While individual case records and practitioner observations demonstrate patient recovery, the lack of systematic documentation, uniform case-taking protocols, institutional research funding, and patient participation in trials hinders mainstream scientific recognition.

This article addresses this critical gap by explicitly examining three specific objectives:

- (1) to describe the symptom pictures of Sepia, Natrum muriaticum, and Pulsatilla and compare them with postpartum depression symptoms.
- (2) to review and evaluate existing research on these remedies while identifying methodological strengths and weaknesses
- (3) to establish clinical safety guidelines prioritizing conventional care.

Furthermore, it serves as a proactive call to action for the medical community to embrace rigorous documentation and research-oriented frameworks.

1.3 Core Objectives of the Study

Primary Objectives of the Investigation:

1. To describe the symptom pictures of Sepia, Natrum muriaticum, and Pulsatilla and compare them with the clinical symptoms of postpartum depression.
2. To review and evaluate existing research on these three remedies in postpartum depression, identifying methodological strengths and weaknesses.
3. To suggest robust safety guidelines for using homeopathic remedies in postpartum depression, ensuring conventional medical care remains the primary clinical priority.

2. Review of Literature

2.1 Epidemiological & Biopsychosocial Determinants of PPD

Recent systematic reviews and umbrella meta-analyses (Ojo-Rowland et al., 2026; Rabiatal Adawiyah et al., 2026; Ji & Wang, 2025) confirm the complex biopsychosocial etiology of PPD. Biological factors include neuroendocrine drops following parturition, gestational diabetes, and cesarean sections.

Psychological factors include antenatal anxiety and depression, while social determinants encompass domestic stress, marital friction, and lack of social support.

2.2 Clinical Screening & Evidence-Based Standard of Care

ACOG (2018; 2023) and NICE (2014) guidelines mandate routine perinatal screening using standardized instruments such as the Edinburgh Postnatal Depression Scale (EPDS) (Cox, Holden, & Sagovsky, 1987) and Hamilton Rating Scale for Depression (HAM-D). Evidence-based psychotherapies (CBT, IPT) and supervised pharmacotherapy represent the gold standard of care.

2.3 Homeopathic Materia Medica & Synthesis of Selected Remedies

Classical materia medica (Hahnemann, 1816; Kent, 1905; Boericke, 1927) details specific constitutional profiles that align with postpartum emotional states.

Table 1: Synthesis of Classical Homeopathic Remedy Profiles & Evidence Base

Remedy & Source	Core Emotional & Mental Profile	Physical Symptoms & Modalities	Key Repertorial Rubrics	Evidence Status & Literature Gap
Sepia officinalis (Cuttlefish Ink)	Profound emotional detachment from infant/spouse; severe irritability; apathy toward maternal duties; emotional burnout.	Pelvic congestion, prolapse/bearing-down sensation. Worse in evening; better with vigorous exercise.	- Mind; indifference to loved ones - Mind; irritability, aversion to child/husband.	Anecdotal / Case Series. Indian journal reports (Sharma 2015) note EPDS improvements. Lacks RCTs.
Pulsatilla nigricans (Windflower)	High emotional dependency; frequent tearfulness; seeking consolation; rapidly changing, volatile moods.	Hormonal fluctuations, digestive upset. Worse in warm, stuffy rooms; better in cool, open air	- Mind; weeping, consolation amels. - Mind; dependency, desire for company	Narrative Case Reports (Gupta 2016). Describes emotional stabilization but lacks systematic controlled trial data
Natrum muriaticum (Sodium Chloride)	Silent, suppressed grief; internal brooding over past loss; weeping in solitude; active aversion to consolation	Recurrent headaches, anemia, mucous dryness. Worse in heat/sun; better in total solitude	- Mind; grief, prolonged - Mind; aversion to consolation	Case Literature Series (Banerjee 2017). Positive bonding narratives; evidence remains unblinded and non-comparative

2.4 Critical Appraisal & The Documentation Challenge in Homeopathic Practice

Systematic reviews (Pilkington et al., 2005; Witt et al., 2013) emphasize that while individual homeopathic case histories demonstrate clinical recovery, the broader medical literature suffers from major methodological constraints: a lack of double-blind RCTs, inconsistent remedy potency selection, and absence of standardized outcome measures.

From the vantage point of clinical practice, homeopathic physicians successfully manage numerous PPD patients by tailoring remedies to individual constitutional types.

However, several compounding factors explain why producing large-scale research papers remains exceptionally difficult.

1. **Absence of Standardized Digital Case-Taking Systems:** Traditional clinical setups often rely on unstructured handwritten records rather than standardized electronic health records (EHR) equipped for clinical trial data extraction.
2. **Time Constraints in High-Volume Clinical Settings:** Practitioners balancing heavy patient loads face severe logistical hurdles in compiling, anonymizing, and tracking longitudinal cohort data for publication.
3. **Patient Participation & Follow-Up Attrition:** Postpartum mothers juggling newborn care frequently experience high attrition rates in multi-month observational studies or clinical trials.
4. **Institutional & Funding Disparities:** Complementary medicine units historically receive limited grant funding for large-scale multi-centric trials compared to conventional pharmaceutical research.

3. Research Methodology

3.1 Design & Review Framework

This study employs structured narrative critical review design, synthesizing multidisciplinary literature from psychiatry, maternal mental health, obstetrics, and classical homeopathic materia medica.

3.2 Inclusion & Exclusion Criteria

- **Inclusion Criteria:** Peer-reviewed empirical studies, clinical guidelines (ACOG, NICE), and authoritative classical texts focusing on perinatal mood disorders within 12 months postpartum.
- **Exclusion Criteria:** Non-perinatal studies, unverified opinion pieces, and unstandardized multi-herbal mixtures lacking classical homeopathic individualization.

4. Discussion, Findings and Clinical Perspectives

4.1 Synthesis of Findings & Clinical Realities

The findings underscore a fundamental dichotomy in modern healthcare: while conventional medicine possesses robust empirical trial frameworks, homeopathic clinical practice relies on individualized qualitative symptom mapping (Sepia, Pulsatilla, Natrum mur) that consistently yields positive patient recovery in daily clinical encounters. However, because practitioners struggle to document and publish these outcomes on a large scale due to structural barriers, homeopathy is frequently mischaracterized as lacking efficacy rather than recognized as lacking institutional research infrastructure.

4.2 Call to Action: Bridging Clinical Practice with Scientific Documentation

To overcome these limitations, the author strongly urges the global homeopathic medical community to enact immediate structural reforms:

- Adoption of Standardized Electronic Case Records (ECRs): Clinics must implement uniform digital case-taking templates capturing baseline EPDS scores, repertorial rubrics, remedy potencies, and longitudinal follow-up metrics.
- Multi-Centric Practice-Based Research Networks (PBRNs): Establishing collaborative networks among homeopathic hospitals and academic institutions will pool clinical data across large patient cohorts.
- Integration into Perinatal Research Grants: Securing institutional support and academic partnerships to fund rigorous, blinded observational and interventional trials.

4.3 Clinical Safety Guidelines & Red-Flag Referral Thresholds

CRITICAL SAFETY PROTOCOL - Emergency Referral Thresholds:

1. Suicidal ideation, self-harm intent, or thoughts of infanticide.
2. Manifestation of psychotic symptoms (hallucinations, persecutory delusions, severe postpartum psychosis).
3. Severe functional incapacitation where the mother cannot care for herself or her infant.
4. Rapid deterioration of mental status despite homeopathic or supportive care .

Mandate: Homeopathic treatment must operate strictly as a supportive adjunct. Under no circumstances should complementary remedies delay immediate psychiatric referral or evidence-based crisis intervention.

5. Conclusion And Future Directions

This article examined PPD and the theoretical descriptions of Sepia, Pulsatilla, and Natrum muriaticum, highlighting the vital role of individualization in homeopathic clinical practice. By candidly addressing the documentation barriers and research limitations that prevent practitioners from publishing large-scale evidence, this paper provides a roadmap for the medical community to elevate homeopathic research standards, ensuring that patient care is fully backed by science.

REFERENCES

1. American College of Obstetricians and Gynecologists. (2018). ACOG Committee Opinion No. 757: Screening for perinatal depression. *Obstetrics & Gynecology*, 132(5),.
2. American College of Obstetricians and Gynecologists. (2023). Screening and diagnosis of mental health conditions during pregnancy and postpartum: Clinical Practice Guideline No. 4. *Obstetrics & Gynecology*, 141(6), 1232–1263
3. Beck, C. T. (2001). Predictors of postpartum depression: An update. *Nursing Research*, 50(5), 275–285
4. Boericke, W. (1927). *Pocket Manual of Homoeopathic Materia Medica* (9th ed.). Boericke & Runyon.
5. Bowlby, J. (1969). *Attachment and Loss: Vol. 1. Attachment*. Basic Books.
6. Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry*, 150(6), 782–786
7. Kent, J. T. (1905). *Lectures on Homoeopathic Materia Medica*. Boericke & Tafel.
8. National Institute for Health and Care Excellence. (2014). Antenatal and postnatal mental health: Clinical management and service guidance (NICE Guideline CG192).
9. O'Hara, M. W., & McCabe, J. E. (2013). Postpartum depression: Current status and future directions. *Annual Review of Clinical Psychology*, 9, 379–407.

10. Ojo-Rowland, M. T., Okobi, O. E., et al. (2026). Postpartum depression in the United States: A systematic review of epidemiology, risk factors, and contemporary management. *Cureus*, 18(5),
11. Patel, V., et al. (2012). Gender, social disadvantage and perinatal depression in low-income countries. *Acta Psychiatrica Scandinavica*, 113(1), 4–11
12. Pilkington, K., Kirkwood, G., Rampes, H., Fisher, P., & Richardson, J. (2005). Homeopathy for depression: A systematic review of the research evidence. *Homeopathy*, 94(3), 153–163.
13. Rabiatal Adawiyah, B. N. I., et al. (2026). Biological, psychological and social determinants of postpartum depression: An umbrella review. *BJPsych Open*, 12(S1), S42
14. Rabiatal Adawiyah, B. N. I., et al. (2026). Biological, psychological and social determinants of postpartum depression: An umbrella review. *BJPsych Open*, 12(S1), S42
15. Sockol, L. E., Epperson, C. N., & Barber, J. P. (2014). Preventing postpartum depression: A meta-analytic review. *Clinical Psychology Review*, 34(5), 389–401
16. Witt, C. M., et al. (2013). Homeopathy for depression: A randomized, partially double-blind, placebo-controlled study. *PLOS ONE*, 8(9),