

Biotechnological Approaches to Biodiversity Conservation and Environmental Management

Khyati Shrivastava¹, Priyanka Tiwari²

^{1,2}Department of Biotechnology, Sam Global University, Bhopal, Madhya Pradesh, India

Abstract

Biodiversity refers to the structural, functional and evolutionary variability among living organisms across genetic, species and ecosystem hierarchical tiers. Accelerating industrialization, habitat fragmentation, deforestation, agrochemical saturation and anthropogenic climate change have initiated a modern mass extinction event that compromises ecological stability and human survival. Traditional conservation frameworks, while foundational, are increasingly constrained by environmental stressors and spatial limitations. Modern biotechnology provides powerful interventions that mitigate biodiversity loss through integrated ex-situ and in-situ conservation strategies. Techniques such as somatic embryogenesis, apical meristem micro propagation, cryopreservation at ultra-low temperatures, synthetic seed technology and high-density DNA banking facilitate the long-term preservation and recovery of endangered, endemic and economically vital germplasm. Recombinant DNA technology, high-throughput molecular marker profiling and next-generation sequencing further allow precise monitoring of population genetics, overcoming bottlenecks such as inbreeding depression. In environmental management, microbial bioremediation, phytoremediation and bio-fermentation processes offer scalable solutions for soil and aquatic detoxification. This review analyses the biotechnological tools applied to biodiversity conservation through a structured literature synthesis and four comparative assessments. Peer-reviewed global assessments show that between 13.6 % (birds) and 40.7 % (amphibians) of assessed species in six major lineages are threatened, with a distinct dominant driver in each lineage, so interventions must be matched to the lineage. An ordinal field-suitability index ranks SSR and SNP platforms highest (4.00 of 5) for conservation genetics, while comparisons of ex-situ and remediation technologies show that they are complementary rather than interchangeable. The review also examines regulatory and bio safety considerations and emerging techniques including genome editing and environmental DNA. The synthesis concludes that biotechnological intervention functions most effectively as a complement to habitat protection rather than as a substitute for it.

Keywords: Conservation biotechnology, Biodiversity loss, Cryopreservation, Micro propagation, DNA banking, Molecular markers, Bio remediation, Genome editing

Highlights

- Ex-situ biotechnological preservation is synthesized across meristem culture, somatic embryogenesis and cryogenic banking, establishing where each method outperforms conventional seed storage.
- Five molecular marker systems are scored on four operational attributes, and a field-suitability index ranks SSR and SNP platforms highest for conservation field work.

- Microbial and plant-based remediation are positioned alongside genomic conservation within a single bio safety and governance framework, showing that biotechnology supplements rather than replaces in-situ protection.




Figure 1. Graphical abstract: integrated biotechnological framework for biodiversity conservation and environmental remediation

1. Introduction

The biosphere is undergoing unprecedented transformation as a result of anthropogenic activity. The term biodiversity, introduced by Walter G. Rosen in the mid-1980s and popularised through the 1986 National Forum on BioDiversity, encompasses the full complex of life forms across all levels of biological organisation [1], [2]. Biodiversity is structurally and functionally stratified into three primary tiers: genetic diversity, the sum of genetic information contained within individual organisms and species populations; species diversity, the variety and abundance of distinct species within defined geographic boundaries; and ecosystem diversity, the dynamic variation of habitats, ecological niches and biophysical processes.

Global biodiversity constitutes the biological foundation of human civilization. It underpins ecosystem services that include atmospheric gas regulation, soil formation and replenishment, nutrient cycling, hydrological filtration, crop pollination and natural pest suppression. Wild germplasm additionally serves as an indispensable reservoir for pharmaceutical research, industrial biomaterials and agricultural breeding programmes directed at food security [3]. Despite these ecological and economic values, global biodiversity is contracting at rates without precedent in human history [2].

The drivers of this crisis are multifaceted and synergistic. Habitat destruction driven by agricultural expansion, urbanization and industrial mining remains the principal cause of terrestrial species extinction. Aquatic biomes face severe degradation from chemical effluents, agricultural run-off, plastic

accumulation and thermal discharge. Anthropogenic climate change is altering temperature and precipitation regimes, forcing species to adapt, to shift their ranges or to face extinction, and it interacts with the other pressures rather than acting independently of them [4]. The consequences are already measurable: amphibians are now the most threatened vertebrate class, with 40.7 % of assessed species threatened, and climate change has become the leading driver of their status deteriorations since 2004 [5]. Invasive alien species, unsustainable overexploitation and emerging infectious pathogens such as chytridiomycosis compound species vulnerability further.

Conventional conservation strategies, centred principally on in-situ protected area networks such as national parks, wildlife sanctuaries and biosphere reserves, and on traditional ex-situ facilities such as botanical gardens, seed banks and zoological collections, remain vital but are increasingly insufficient when operated in isolation. In-situ reserves are vulnerable to external pressures including climate shift, poaching, edge effects and catastrophic habitat destruction. Traditional seed banks face constraints when managing recalcitrant or vegetatively propagated species whose seeds tolerate neither desiccation nor freezing [6]. Modern biotechnology offers a set of tools that supplement and extend these classical conservation paradigms rather than displacing them [7].

1.1 Importance of the Study

The importance of this study rests on three considerations. First, the pace of loss now exceeds what habitat protection alone can reverse. Many populations have already fallen below the size at which demographic stochasticity and inbreeding depression drive further decline irrespective of habitat status, and recovering them requires genetic management tools that only biotechnology can provide [8]. Second, the relevant literature is fragmented across plant tissue culture, cryobiology, conservation genetics and environmental microbiology. Conservation managers, particularly in biodiversity-rich but resource-limited countries such as India, therefore lack a single reference that compares the available tools on common criteria. Third, the choice of tool has direct consequences for cost and outcome: an unsuitable storage protocol or marker platform can waste irreplaceable collections. A structured comparative synthesis that links threat profiles to matched interventions therefore has immediate practical value for germplasm banks, botanical gardens, zoological institutions and pollution-control agencies.

1.2 Objectives of the Study

This review provides a structured analysis of the intersection between biotechnology, biodiversity preservation and environmental management. The specific objectives of the study are as follows:

- To identify and analyse the primary anthropogenic drivers accelerating biological extinction across major taxonomic groups.
- To review the methodology of advanced ex-situ propagation and preservation technologies, including tissue culture, micro propagation, cryopreservation and synthetic seed production.
- To evaluate the utility of molecular markers, genomic profiling and high-throughput sequencing in analysing genetic diversity and preventing inbreeding bottlenecks in threatened populations.
- To assess environmental biotechnology solutions, with emphasis on microbial bioremediation and phytoremediation for the restoration of polluted ecosystems.
- To examine future trajectories, including genome editing for climate resilience, synthetic biology and the international regulatory frameworks governing biosafety.

1.3 Research Contributions

The principal contributions of this review are as follows:

- An integrated procedural framework (Figure 2) that places plant tissue culture, cryopreservation, molecular marker profiling and bioremediation within a single four-workflow structure, with protocol parameters stated explicitly.
- A taxon-level synthesis of peer-reviewed extinction-risk assessments for six major lineages (Table 1, Figure 3) that matches the dominant threat in each lineage to the biotechnological intervention best suited to it.
- A comparative evaluation of four ex-situ preservation technologies (Table 2) by material conserved, storage period, infrastructure requirement and principal limitation.
- A transparent ordinal scoring scheme and field-suitability index for five molecular marker systems (Equation (2), Table 3, Figure 4), which gives conservation managers an explicit basis for platform selection.
- A comparison of in-situ, ex-situ and plant-based remediation routes (Table 4), positioned within the biosafety and governance framework under which all of these tools must operate.

1.4 Organization of the Article

This article is organised into five sections. Section 1 introduces the concept of biodiversity, the drivers of its decline, and the importance, objectives and contributions of the study. Section 2 reviews the related literature spanning ethnobotanical survey, in-vitro conservation, cryogenic storage, DNA banking, molecular marker development and bioremediation, and identifies the gaps that this review addresses. Section 3 describes the review design, the literature search strategy and the assessment metrics, and then sets out the procedural protocols that govern biotechnological conservation practice. Section 4 presents the results as four comparative assessments and discusses the efficacy, limitations and bio safety implications of these tools relative to traditional methods. Section 5 concludes the review and outlines the future scope for research.

2. Related Work

The evolution of conservation biotechnology reflects several decades of interdisciplinary research bridging molecular biology, ecology, forestry and environmental engineering. A synthesis of the seminal and contemporary literature identifies the milestones summarized below.

Ethnobotanical survey documented the medicinal uses of plants in the degraded Chandaka forest patches of Bhubaneswar in eastern India and the dependence of local communities on these species [9]. Such dependence places sustained harvesting pressure on wild stands and underlines the need for micro propagation pipelines that can supply medicinal flora without further depleting natural populations.

A decade of progress in the in-vitro conservation of threatened plants and showed that tissue culture had become a routine component of conservation programmed at botanic gardens [10]. More recently, the surveyed micro propagation techniques for rare plant species and emphasised the importance of explant choice, growth-regulator balance and genetic-stability testing of regenerated plants [11].

In the domain of cryogenic storage and recalcitrant germplasm, evaluated in-vitro culture combined with liquid nitrogen storage for endangered Australian flora [12]. Their work established that cryogenic storage offers effectively indefinite viability for species whose seeds cannot be held in conventional seed banks, a conclusion reinforced subsequently identified the practical barriers to implementing shoot-tip cryopreservation at genebank scale, including species-specific protocol optimization, staff training and the cost of maintaining a liquid nitrogen supply [13].

It assessed biotechnological applications across crop, forest, animal and aquatic genetic resources within the framework of the Food and Agriculture Organization, emphasising how molecular characterisation, cryopreservation and in-vitro storage safeguard crop wild relatives and livestock breeds so that climate-resilient traits remain available to future breeding programmes. A blueprint for DNA banking, illustrating that long-term storage of purified genomic DNA offers a stable, compact and cost-effective backup for biodiversity research, enabling subsequent analysis even where source populations have been lost [14].

The development of molecular tools was reviewed who argued that the value of DNA-level data depends critically on the analytical framework applied to it [15]. This argument into the genomics era, showing that genome-wide data now provide precise estimates of effective population size, inbreeding and adaptive variation that directly inform captive breeding and reintroduction. The reviewed environmental DNA, which permits species detection from water, soil and sediment samples without capturing organisms, while cautioning about imperfect detection and the difficulty of estimating abundance [16].

Bioremediation techniques have been classified according to their site of application, and their underlying principles, advantages, and limitations have been compared for environmental remediation. The mechanisms through which plants extract, stabilise, and sequester heavy metals from contaminated soils have also been reviewed. Additionally, current strategies for plant biodiversity conservation and the international conventions governing these efforts have been discussed [17]. The potential applications and associated risks of synthetic biology, including genome-editing technologies, in mitigating global biodiversity loss have also been explored.

Taken together, the literature confirms the transformative potential of biotechnology in conservation. Three gaps nonetheless persist. The first concerns the integration of genomic profiling with operational field management, which remains weakly developed relative to the maturity of the underlying sequencing technology. The second concerns the scaling of bioremediation from controlled trials to complex, mixed chemical pollution in the field[18]. The third is the absence of explicit, reproducible criteria for choosing among tools, since most reviews describe techniques individually rather than comparing them on common attributes. This review addresses these gaps by presenting the available tools within a single integrated framework and by comparing them on stated criteria.

3. Materials and Methods

This study is a structured narrative review supported by a comparative synthesis of secondary data. The methodology comprised four stages: (i) identification and screening of the literature; (ii) extraction of quantitative and descriptive data; (iii) derivation of comparative metrics; and (iv) consolidation of standard laboratory and field protocols into an integrated procedural framework (Figure 2). No new primary experimental data were generated for this review.

3.1 Review Design and Literature Search Strategy

Peer-reviewed literature was identified through searches of Scopus, Web of Science, Pub Med and Google Scholar, supplemented by backward citation tracking from key reviews. Search strings combined the terms conservation biotechnology, micro propagation, cryopreservation, synthetic seed, DNA banking, molecular marker, population genomics, environmental DNA, bioremediation and phytoremediation with biodiversity, threatened species or endangered [19]. Only journal articles, books and conference proceedings were retained; web pages and other grey literature were excluded. Studies

were included where they (a) described a biotechnological method applied to the conservation of wild species or genetic resources, or to the remediation of contaminated environments, (b) reported protocol parameters or comparative performance, or (c) reported a global, taxon-wide extinction-risk assessment. Priority was given to global assessments and to syntheses published since 2020 so that the comparative data reflect current conditions.

3.2 Data Extraction and Assessment Metrics

For each taxonomic group, the number of species assessed, the proportion threatened and the dominant extinction drivers were extracted from the most recent peer-reviewed global assessment available: amphibians [5], mammals [20], gymnosperms [21], birds [22], reptiles [23] and reef-building corals [24]. The proportion of species threatened follows the IUCN Red List convention, in which species listed as Critically Endangered (CR), Endangered (EN) or Vulnerable (VU) are counted as threatened, and Extinct (EX) and Data Deficient (DD) species are excluded from the denominator, as expressed in Equation (1).

$$T = \frac{N_{CR} + N_{EN} + N_{VU}}{N_A - N_{EX} - N_{DD}} \times 100 \quad (1)$$

where T is the percentage of species threatened and N_A is the total number of species assessed. Because the cited assessments were conducted in different years, the values are compared as relative indicators of threat rather than as a synchronous census.

Marker system characteristics were compiled from the conservation genetics literature [8], [14], [15] and described on five operational attributes: genomic abundance, degree of polymorphism, mode of inheritance, DNA quality requirement and principal conservation application (Table 3). To convert these descriptive categories into a comparable measure, the first four attributes were coded on an ordinal scale of 1 to 5, where 5 denotes the condition most favourable to conservation field work. Abundance and polymorphism were coded Low = 1, Medium = 2, High = 3, Very high = 4 and Ultra high = 5. Inheritance was coded Dominant = 1 and Co-dominant = 5, because only co-dominant markers distinguish heterozygotes, which pedigree and inbreeding analysis requires. DNA quality requirement was reverse-coded as tolerance of degraded DNA, from High (pure, undegraded) = 1, High = 2, Medium to high = 3 and Medium = 4 to Low to medium = 5. A field-suitability index (FSI) was then computed for each marker system as the unweighted mean of its attribute scores, as given in Equation (2).

$$FSI_m = \frac{1}{k} \sum_{j=1}^k s_{mj} \quad (2)$$

where s_{mj} is the ordinal score of marker system m on attribute j, and k = 4 is the number of attributes scored. Equal weighting was adopted deliberately so that the index remains transparent and reproducible; users with specific objectives may re-weight the attributes accordingly. Ex-situ preservation technologies and bioremediation approaches were compared qualitatively on material conserved or treated, time scale, infrastructure requirement and principal limitation, using the sources cited in Tables 2 and 4.

3.3 Plant Tissue Culture and Micro propagation

Plant tissue culture relies on totipotency, the capacity of a single plant cell to divide and differentiate into a complete organism. For threatened flora, micro propagation provides a rapid route to pathogen-free clonal offspring [20], [21] through the following sequence.

1. Explant selection and preparation. Active meristematic tissues, apical buds or nodal segments are collected from healthy mother plants. Tissues are pre-washed under running tap water for approximately 30 minutes with gentle detergent treatment.
2. Surface sterilization. Explants are transferred to a laminar airflow cabinet and treated with 70 % ethanol for 60 seconds, followed by immersion in 1.5 % sodium hypochlorite or 0.1 % mercuric chloride with Tween-20 for 8 to 12 minutes. Explants are then rinsed five times in sterile distilled water.
3. Culture medium formulation. Explants are inoculated onto Murashige and Skoog (MS) basal medium containing 3 % sucrose and 0.8 % agar, adjusted to pH 5.7 to 5.8. Growth regulators, comprising auxins such as indole-3-acetic acid (IAA) or naphthaleneacetic acid (NAA) and cytokinins such as 6-benzylaminopurine (BAP) or kinetin, are added in controlled ratios to promote shoot multiplication or callus formation.
4. Incubation conditions. Cultures are maintained in controlled growth chambers at 25 ± 2 °C and 60 to 70 % relative humidity, under a 16-hour photoperiod provided by cool-white fluorescent lamps at approximately 3000 lux.
5. Rooting and acclimatisation. Elongated shoots are transferred to half-strength MS medium supplemented with indole-3-butyric acid to induce rooting. Rooted plantlets are moved to hardening pots containing a sterile 1:1 mixture of vermiculite and soilrite, acclimatised in shade houses and subsequently reintroduced to their natural habitat.

3.4 Cryopreservation and Synthetic Seed Technology

Cryopreservation stores biological tissues such as meristems, embryos, pollen and cell suspensions at ultra-low temperature, in liquid nitrogen at -196 °C or in its vapour phase at approximately -150 °C. At these temperatures enzymatic degradation and metabolic activity effectively cease, permitting storage over indefinite periods [19].

Before cryoprotection, explants are commonly pre-cultured on sucrose-enriched medium to increase their tolerance of dehydration [13]. To prevent lethal intracellular ice crystal formation during cooling, tissues are then treated with cryoprotectant formulations such as Plant Vitrification Solution 2 (PVS2), comprising 30 % (w/v) glycerol, 15 % (w/v) ethylene glycol and 15 % (w/v) dimethyl sulfoxide in liquid medium containing 0.4 M sucrose. Droplet-vitrification protocols apply the solution to explants held on aluminium foil strips before direct immersion in liquid nitrogen. On rewarming, the foil strips are transferred directly into a 1.2 M sucrose unloading solution, whereas samples vitrified in cryovials are first rewarmed in a water bath at approximately 40 °C; explants are then washed and transferred to recovery medium.

For animal germplasm, cryopreservation is applied to spermatozoa, oocytes, embryos and somatic cell lines. Assisted reproductive technologies based on cryopreserved sperm have produced offspring in several threatened amphibian species [25], and cryopreserved coral sperm has been used to cross geographically separated populations of a critically endangered coral, providing a practical route to assisted gene flow for reef restoration [26].

Synthetic seed technology complements cryopreservation. Somatic embryos or nodal explants are encapsulated in a 3 % sodium alginate matrix and dropped into a 100 mM calcium chloride solution to form protective calcium-alginate beads. These artificial seeds can be stored for short periods, transported

and germinated directly under sterile conditions, which reduces the logistical burden of moving living material between collection sites and laboratories [6].

3.5 DNA Banking and Molecular Marker Diagnostics

DNA banking involves the systematic extraction, purification and cold storage at -80°C of genomic DNA from endangered flora and fauna [14]. Genomic DNA is extracted using modified cetyltrimethylammonium bromide (CTAB) protocols for plant tissue, or proteinase K digestion with phenol-chloroform extraction for animal tissue. Molecular markers are then applied to evaluate genetic diversity, population structure and gene flow [15]. The five platforms compared in Section 4 are characterised below.

- Restriction fragment length polymorphisms (RFLP) are co-dominant, hybridisation-based markers that require large quantities of pure, undegraded DNA and are now used mainly for historical phylogenetic comparison.
- Random amplified polymorphic DNA (RAPD) markers use short arbitrary primers, need no prior sequence information and tolerate low-quality DNA, but are dominant and have limited reproducibility between laboratories.
- Amplified fragment length polymorphisms (AFLP) provide a high-throughput dominant marker technique giving genome-wide coverage without prior sequence knowledge.
- Simple sequence repeats (SSR), or microsatellites, are co-dominant, highly polymorphic loci used to detect subtle genetic differences between isolated populations and to assess rates of inbreeding.
- Single nucleotide polymorphisms (SNP) analysed by next-generation sequencing (NGS) evaluate functional variation, adaptive traits and evolutionary history across many thousands of loci simultaneously [8].
- Environmental DNA (e DNA) metabarcoding amplifies DNA shed by organisms into water, soil or sediment, allowing many species to be detected from a single environmental sample without capture [16].

3.6 Bio remediation and Environmental Detoxification

Environmental biotechnology employs biological organisms, including bacteria, fungi, algae and higher plants, to degrade, neutralise or sequester environmental pollutants [17]. Three procedural families are distinguished.

- In-situ bioremediation includes biosparging, bioventing and bioaugmentation, in which specialised microbial strains such as *Pseudomonas putida* and *Rhodococcus erythropolis* are introduced directly into contaminated soil or groundwater to metabolise hydrocarbons and organic solvents.
- Ex-situ bioremediation excavates contaminated soil or wastewater for processing in slurry-phase bioreactors, biopiles, landfarming units or composting beds under controlled oxygen and nutrient supply [17].
- Phytoremediation employs hyperaccumulating plant species such as *Brassica juncea* and *Helianthus annuus* to absorb heavy metals including cadmium, lead, arsenic and chromium from soil through phytoextraction, phytostabilisation or rhizofiltration [18].

3.7 Integrated Procedural Framework

Figure 2 summarises the four workflows described in Sections 3.3 to 3.6 as sequential pipelines. Each row begins with the conservation objective on the left and proceeds through its principal procedural stages. Workflows 1 and 2 are linked, because in-vitro cultures supply the explants for cryopreservation,

and workflow 3 provides the genetic monitoring that verifies the fidelity of material regenerated from workflows 1 and 2 before reintroduction. Workflow 4 operates on the habitat rather than on the organism, restoring the sites to which conserved material can ultimately be returned.




Figure 2. Integrated conservation biotechnology procedural framework: the four standardized workflows described in Sections 3.3 to 3.6

4. Results and Discussion

4.1 Results

4.1.1 Global Assessment of Biodiversity Threats

Table 1 summarises the extinction-risk profile of six major lineages, compiled from the peer-reviewed global assessments listed in Section 3.2, together with the biotechnological interventions most closely matched to each profile. The proportions threatened are plotted in Figure 3.

Table 1. Global extinction vulnerability across six major lineages and the matched biotechnological intervention

Taxonomic Group	Species Assessed	Threatened (%)	Principal Extinction Drivers	Matched Biotechnological Intervention	Source
Amphibians	8,011	40.7	Chytrid fungal disease, habitat loss, climate change	Sperm cryopreservation and assisted reproduction; biobanking of cell lines	[5], [25]

Taxonomic Group	Species Assessed	Threatened (%)	Principal Extinction Drivers	Matched Biotechnological Intervention	Source
Gymnosperms	1,004	39.9	Deforestation, logging, habitat loss in restricted ranges	Meristem micropropagation; somatic embryogenesis; shoot-tip cryopreservation	[6], [21]
Reef-building corals	845	32.8 ^a	Bleaching and disease driven by elevated sea temperature; local impacts	Sperm cryopreservation for assisted gene flow; heat-evolved symbionts	[24], [26], [27]
Mammals	5,487	25.4 ^b	Habitat loss and degradation, harvesting	Genome resource banking; molecular monitoring of captive populations	[3], [20], [23]
Reptiles	10,196	21.1	Agriculture, logging, urban development, invasive species	Assisted reproduction; molecular sexing and population markers	[23], [25]
Birds	≈11,000	13.6 ^b	Habitat loss, climate change, overexploitation, invasive species	Genetic profiling; genomic monitoring of small populations	[22], [23]

^a Proportion of the 704 coral species with sufficient data for assignment of a conservation status [24]. ^b Threatened proportion as reported by Cox et al. [23]; species-assessed totals from [20] (mammals) and [22] (birds).

Three results emerge from Table 1 and Figure 3. First, the proportion of species threatened varies about threefold across lineages, from 13.6 % in birds to 40.7 % in amphibians, with a mean of 28.9 % across the six groups. Second, the two most threatened groups, amphibians and gymnosperms, are phylogenetically and ecologically unrelated, which indicates that high threat is not confined to any single branch of the tree of life. Third, the dominant driver differs between groups: disease and climate change for amphibians, forest loss within restricted ranges for gymnosperms and thermal stress for corals, whereas habitat loss and overexploitation dominate for mammals, reptiles and birds.




Figure 3. Proportion of assessed species threatened with extinction (CR + EN + VU) in six major lineages; n = number of species assessed. Data compiled from Table 1

In Figure 3 the lineages are ordered by threat level. Each bar gives the percentage of species classified as CR, EN or VU, and the label inside each bar gives the number of species assessed. The figure shows that proportional threat is not a function of assessment effort: reptiles and birds, the two largest samples, show the lowest proportions threatened, whereas the much smaller gymnosperm sample ranks second. The practical consequence is that biotechnological intervention must be matched to the lineage rather than applied uniformly. Gamete cryopreservation addresses amphibian and coral decline because it preserves genetic diversity independently of the disease- or heat-affected habitat, whereas meristem culture and shoot-tip cryopreservation suit gymnosperms, whose long generation times slow recovery from seed and for which clonal propagation preserves specific genotypes.

4.1.2 Comparative Evaluation of Ex-situ Preservation Technologies

Table 2 compares the four principal ex-situ technologies on the material they conserve, the storage period they support, their infrastructure requirements and their chief limitation.

Table 2. Comparative evaluation of ex-situ preservation technologies used in conservation biotechnology

Technology	Material Conserved	Storage Period	Infrastructure Requirement	Principal Limitation	Source
Micropropagation and slow-growth storage	Shoots, nodal segments, somatic embryos	Months to a few years between subcultures	Laminar cabinets, airflow growth trained rooms, staff	Somaclonal variation and contamination risk over repeated subculture	[10], [11]

Technology	Material Conserved	Storage Period	Infrastructure Requirement	Principal Limitation	Source
Cryopreservation	Shoot tips, embryos, pollen, gametes, cell lines	Long-term; theoretically indefinite	Reliable liquid nitrogen supply, cryotanks, species-specific protocols	Protocol optimisation needed for each species; recovery rates vary	[6], [13]
Synthetic seeds	Encapsulated somatic embryos or nodal explants	Short to medium term unless combined with cryopreservation	Basic tissue culture facilities; sodium alginate and calcium chloride	Limited storage life; conversion to plantlets varies between species	[6]
DNA banking	Purified genomic DNA	Long-term at -80°C or below	Ultra-low freezers; extraction and quality-control facilities	Preserves genetic information only; cannot regenerate living organisms	[14]

Table 2 shows that the four technologies are complementary rather than interchangeable. Micro propagation provides the living material and the multiplication capacity on which the other methods depend, but it is the least suitable for long-term storage because every subculture adds risk of somaclonal variation and contamination. Cryopreservation removes that risk and offers the longest storage period, at the cost of a continuous liquid nitrogen supply and species-specific protocol development [13]. Synthetic seeds occupy an intermediate position and are useful mainly for exchange and short-term holding of material between sites. DNA banking is the most compact option but preserves information rather than organisms. An effective germplasm programme therefore combines at least two of these technologies: a living collection maintained in vitro and a cryopreserved or DNA-banked back-up.

4.1.3 Comparative Evaluation of Molecular Marker Systems

Selection of an appropriate molecular marker system is decisive for population structure analysis and for guiding reintroduction programmes. Table 3 summarises the operational characteristics of the five marker platforms described in Section 3.5, together with the field-suitability index calculated from Equation (2).

Table 3. Operational comparison of molecular marker technologies used in conservation genetics, with field-suitability index (FSI)

Marker System	Abundance	Polymorphism	Inheritance	DNA Quality Needed	Primary Conservation Application	FSI
RFLP	Low	Medium	Co-dominant	High (pure, undegraded)	Historical phylogenetic studies	2.25

Marker System	Abundance	Polymorphism	Inheritance	DNA Quality Needed	Primary Conservation Application	FSI
RAPD	High	High	Dominant	Low to medium	Initial genetic screening of wild flora	3.00
AFLP	Very high	High	Dominant	High	Genome mapping and species identification	2.50
SSR (microsatellite)	High	Very high	Co-dominant	Medium	Inbreeding and pedigree analysis	4.00
SNP / NGS	Ultra high	High	Co-dominant	Medium to high	Genome-wide conservation adaptation studies	4.00

The comparison shows a consistent trade-off. Platforms requiring high-purity, undegraded DNA deliver clean phylogenetic resolution but are poorly suited to the degraded material typically recovered from non-invasive field sampling such as hair, faeces or shed skin, whereas microsatellite and SNP platforms tolerate intermediate DNA quality while retaining co-dominant inheritance, which is the property required for pedigree and inbreeding analysis [8].




Figure 4. Heat map of ordinal scores for five molecular marker systems on four operational attributes, with the resulting field-suitability index (FSI, Equation (2)). Scores derived from Table

3

Figure 4 displays the ordinal scores behind the index as a heat map, in which darker cells indicate conditions more favourable to field conservation work; the column to the right of the red divider gives the FSI. SSR and SNP/NGS platforms share the highest index (4.00), but for different reasons: SSR

markers combine the highest polymorphism with good tolerance of degraded DNA, whereas SNP/NGS platforms derive their score from genome-wide abundance. RAPD scores 3.00 on the strength of its tolerance of low-quality DNA, but its dominant inheritance and limited reproducibility restrict it to preliminary screening [15]. AFLP (2.50) and RFLP (2.25) score lowest because both require high-quality DNA that is rarely obtainable from field samples.

4.1.4 Comparative Evaluation of Bio remediation Approaches

Table 4 compares the three bioremediation routes described in Section 3.6 on their mode of action, the contaminants they address, and their main advantage and limitation.

Table 4. Comparative evaluation of bioremediation and phytoremediation approaches for habitat restoration

Approach	Mode of Action	Typical Contaminants	Main Advantage	Principal Limitation	Source
In-situ bioremediation (bioventing, biosparging, bioaugmentation)	Microbial degradation at the contaminated site, enhanced by aeration or introduced strains	Petroleum hydrocarbons, organic solvents	No excavation; lower cost; minimal disturbance of soil structure	Subsurface is difficult to monitor and control; treatment is slower	[17]
Ex-situ bioremediation (biopiles, landfarming, slurry-phase bioreactors)	Excavated soil or water treated under controlled conditions	Hydrocarbons, pesticides and other organic pollutants	Close control of oxygen, nutrients and temperature; faster treatment	Excavation and transport costs; disturbance of the site	[17]
Phytoremediation (phytoextraction, phytostabilisation, rhizofiltration)	Plants absorb, immobilise or filter contaminants	Heavy metals (Cd, Pb, As, Cr) and some organic pollutants	Low cost and solar-driven; improves soil cover and landscape	Slow; limited to rooting depth; contaminated biomass requires safe disposal	[18]

Table 4 indicates that the choice among remediation routes is governed by the contaminant and by the value placed on soil integrity. Where organic pollutants must be removed quickly and the site can tolerate disturbance, ex-situ bioreactors offer the greatest process control. Where soil structure and its associated biodiversity are themselves conservation targets, in-situ approaches and phytoremediation are preferable despite slower rates, because they leave the soil profile intact [17]. For heavy metals, which cannot be degraded, phytoextraction physically removes the metal from the soil into harvestable biomass, whereas microbial processes can only alter its mobility or chemical form; the harvested biomass must then be handled as hazardous waste [18].

4.1.5 Summary of Key Findings

The results of the four comparative assessments can be briefly reviewed as follows:

- Threat is high and uneven: between 13.6 % and 40.7 % of assessed species are threatened across the six lineages examined (mean 28.9 %), and the dominant driver differs between lineages.
- Ex-situ technologies are complementary: micropropagation supplies material, cryopreservation provides the longest secure storage, and DNA banking offers a compact genetic back-up.
- SSR and SNP/NGS platforms have the highest field-suitability index (4.00 of 5), whereas RFLP (2.25) and AFLP (2.50) are least suited to degraded field samples.
- The choice of remediation route depends on the contaminant type and on whether soil integrity is itself a conservation objective.

4.2 Discussion

The integration of biotechnological techniques into conservation biology addresses weaknesses inherent to traditional conservation methods. In-situ reserves, while critical for protecting intact ecosystems, are increasingly exposed to shifting climate conditions that can move suitable habitat outside reserve boundaries [4], as well as to illegal land conversion and localised environmental disaster. Where a wild population falls below a critical threshold, demographic stochasticity and inbreeding depression can accelerate extinction irrespective of physical habitat protection, which is precisely the failure mode that the marker platforms compared in Table 3 are designed to detect before it becomes irreversible [8].

Ex-situ biotechnological methods provide an essential safeguard against that failure mode. Micropropagation permits rapid multiplication of critically endangered plants from minimal initial tissue, circumventing seed dormancy and poor germination rates [11]. Cryopreservation provides near-indefinite storage without the somaclonal variation or genetic drift associated with continuous in-vitro subculture [6], [12]. DNA banking preserves genomic information even where wild populations have been extirpated [14]. The taxon-specific matching set out in Table 1 matters here: gamete cryopreservation addresses amphibian and coral decline in a way that micropropagation cannot, while meristem culture addresses gymnosperm loss in a way that gamete banking cannot.

In pollution management, bioremediation and phytoremediation offer sustainable alternatives to conventional physical and chemical clean-up. Excavation and incineration frequently cause severe secondary damage, destroying soil structure and soil fauna. By contrast, bioaugmentation using adapted microbial consortia breaks down toxic aromatic hydrocarbons, synthetic pesticides and industrial dyes into less harmful products, principally carbon dioxide, water and biomass [17]. Hyperaccumulating plants remove heavy metals from soil, restoring fertility and reducing toxic run-off into aquatic ecosystems [18].

Technical and biosafety constraints must nonetheless be managed. In-vitro micro propagation carries a risk of somaclonal variation where growth regulator concentrations are improperly balanced [10]. Cryopreservation requires specialised equipment, a reliable liquid nitrogen supply chain and species-specific protocol optimisation, which limits deployment in exactly those regions where endemic diversity is highest [13]. The release of genetically engineered organisms or engineered biocontrol agents into wild ecosystems raises further concerns regarding self-dissemination, gene transfer to wild relatives and non-target ecological effects [19].

Deployment must therefore be governed by robust biosafety protocols and legal frameworks. The Convention on Biological Diversity provides the overarching framework for the conservation and sustainable use of biodiversity [2], and its Cartagena Protocol on Biosafety sets international standards for the transboundary movement and management of living modified organisms. Rigorous multi-tier risk

assessment should precede any field introduction of genetically modified biocontrol agents or transgenic flora, in order to prevent unintended disruption of wild food webs. Compliance with the Nagoya Protocol ensures that genetic resources and traditional knowledge accessed for biotechnological research are used with fair and equitable benefit-sharing among indigenous and local communities [19]. Public concern surrounding genomic modification, de-extinction and synthetic biology requires transparent communication, functioning ethical review committees and sustained interdisciplinary collaboration.

Read together, the four comparative assessments point to a single operational sequence. Threat assessment (Table 1) identifies which lineages require intervention and what the dominant driver is; the ex-situ comparison (Table 2) determines how material should be secured; marker profiling (Table 3, Figure 4) verifies that the secured material is genetically representative and monitors the populations to which it is returned; and remediation (Table 4) restores the habitat that reintroduction requires. Omitting any stage weakens the others, which is the principal practical argument for the integrated framework presented in Figure 2.

Emerging tools are extending this sequence. Environmental DNA surveys allow managers to confirm the presence of rare species and to monitor reintroduced populations without capture [16], and laboratory evolution of coral endosymbionts has already produced symbiont strains that raise the bleaching tolerance of coral larvae [27]. Synthetic biology and genome editing may in future supply similar adaptive capacity to other lineages, but they raise questions of self-dissemination, gene flow to wild relatives and public acceptance that remain unresolved [19].

The limitations of the present analysis should be acknowledged. The comparative tables rest on secondary data drawn from published assessments rather than on primary measurement, and those assessments were conducted between 2008 and 2023 using differing survey effort, so the threatened proportions in Table 1 are indicative rather than strictly comparable. Plants and invertebrates remain substantially under-assessed relative to vertebrates. The field-suitability index uses a coarse ordinal scale with equal weights, and a different weighting would change the ranking of intermediate platforms such as RAPD and AFLP. Direct empirical field trials of the micro propagation and bioremediation protocols described in Section 3 were beyond the scope of this review, and reported efficacy therefore reflects the conditions of the originating studies. Future work should prioritise controlled comparison of protocol variants across a common species set, which would allow the qualitative rankings presented here to be replaced by quantitative effect estimates.

5. Conclusion and Future Scope

Biotechnology represents a vital frontier in the effort to slow biodiversity loss and to mitigate environmental degradation. This review has shown that tissue culture micropropagation, ultra-low temperature cryopreservation, synthetic seed technology, DNA banking and molecular marker diagnostics together allow genetic diversity to be secured across endangered flora and fauna, including the recalcitrant and vegetatively propagated species that conventional seed banking cannot hold. Comparative assessment of extinction risk across six major lineages showed that between 13.6 % and 40.7 % of assessed species are threatened and that the dominant driver differs between lineages, so intervention must be matched to the lineage rather than applied uniformly, with gamete cryopreservation suited to amphibian and coral decline and meristem culture suited to gymnosperm loss. Comparison of ex-situ technologies showed that they are complementary, and that a secure programme combines a living in-vitro collection with a cryopreserved or DNA-banked back-up. The field-suitability index

identified SSR and SNP/NGS platforms (FSI = 4.00) as best suited to field conservation genetics, providing conservation managers with an explicit, reproducible basis for platform selection. In parallel, microbial bioremediation and phytoremediation offer scalable and comparatively non-destructive routes to the restoration of contaminated soil and water systems. The principal limitation of these approaches is that they remain resource-intensive and species-specific, requiring specialised infrastructure and reliable supply chains that are frequently least available in the regions of highest endemic diversity. The central recommendation arising from this synthesis is that biotechnological intervention be positioned as a complement to habitat protection rather than as a substitute for it, since preserved germplasm has conservation value only where a habitat remains to receive it.

Several developments are likely to shape the future scope of this field. Precise genome editing offers a route to introducing climate-adaptive traits, such as fungal disease resistance in threatened forest trees, without the introduction of foreign DNA, although its release into the wild requires careful governance [19]. High-throughput sequencing of environmental DNA recovered from soil, water and air permits non-invasive monitoring of elusive species and microbial communities at landscape scale, and represents the most plausible near-term route to closing the gap between genomic capability and operational field management identified in Section 2 [16]. Assisted evolution of symbionts, already demonstrated for corals, may be extended to other holobionts under thermal stress [27]. Advanced reproductive technologies combining somatic cell nuclear transfer, induced pluripotent stem cells and primordial germ cell manipulation offer theoretical pathways for restoring functionally extinct species, though these remain experimental and ethically contested. Finally, the establishment of automated, remotely monitored and internationally networked cryogenic biobanks would standardise conservation protocols and streamline germplasm exchange between countries. Research directed at these areas, coupled with field validation of existing protocols and refinement of the comparative indices proposed here, will determine whether conservation biotechnology fulfils its demonstrated potential.

References

1. E.O. Wilson, "BioDiversity," National Academy Press, USA, pp.1-521, 1988.
2. Amanjot Kaur, "Conservation of Plant Biodiversity: Current Strategies and Future Needs," International Journal of Scientific Research in Biological Sciences, Vol.5, Issue.4, pp.109-113, 2018.
3. P. Lidder, A. Sonnino, "Biotechnologies for the Management of Genetic Resources for Food and Agriculture," Advances in Genetics, Vol.78, pp.1-167, 2012.
4. C. Bellard, C. Bertelsmeier, P. Leadley, W. Thuiller, F. Courchamp, "Impacts of Climate Change on the Future of Biodiversity," Ecology Letters, Vol.15, Issue.4, pp.365-377, 2012.
5. J.A. Luedtke, J. Chanson, K. Neam, L. Hobin, A.O. Maciel, A. Catenazzi, et al., "Ongoing Declines for the World's Amphibians in the Face of Emerging Threats," Nature, Vol.622, Issue.7982, pp.308-314, 2023.
6. F. Engelmann, "Use of Biotechnologies for the Conservation of Plant Biodiversity," In Vitro Cellular & Developmental Biology – Plant, Vol.47, Issue.1, pp.5-16, 2011.
7. B.M. Reed, V. Sarasan, M. Kane, E. Bunn, V.C. Pence, "Biodiversity Conservation and Conservation Biotechnology Tools," In Vitro Cellular & Developmental Biology – Plant, Vol.47, Issue.1, pp.1-4, 2011.
8. P.A. Hohenlohe, W.C. Funk, O.P. Rajora, "Population Genomics for Wildlife Conservation and Management," Molecular Ecology, Vol.30, Issue.1, pp.62-82, 2021.

9. K.K. Behera, S. Sahoo, S. Patra, “Floristic and Medicinal Uses of Some Plants of Chandaka Denudated Forest Patches of Bhubaneswar, Orissa, India,” *Ethnobotanical Leaflets*, Vol.12, pp.1043-1053, 2008.
10. V. Sarasan, R. Cripps, M.M. Ramsay, C. Atherton, M. McMichen, G. Prendergast, J.K. Rowntree, “Conservation In Vitro of Threatened Plants – Progress in the Past Decade,” *In Vitro Cellular & Developmental Biology – Plant*, Vol.42, Issue.3, pp.206-214, 2006.
11. V.A. Chokheli, P.A. Dmitriev, V.D. Rajput, S.D. Bakulin, A.S. Azarov, T.V. Varduni, et al., “Recent Development in Micropropagation Techniques for Rare Plant Species,” *Plants*, Vol.9, Issue.12, Article 1733, 2020.
12. E. Bunn, S.R. Turner, M. Panaia, K.W. Dixon, “The Contribution of In Vitro Technology and Cryogenic Storage to Conservation of Indigenous Plants,” *Australian Journal of Botany*, Vol.55, Issue.3, pp.345-355, 2007.
13. J.C. Bettoni, R. Bonnart, G.M. Volk, “Challenges in Implementing Plant Shoot Tip Cryopreservation Technologies,” *Plant Cell, Tissue and Organ Culture*, Vol.144, Issue.1, pp.21-34, 2021.
14. T.R. Hodkinson, S. Waldren, J.A.N. Parnell, C.T. Kelleher, K. Salamin, N. Salamin, “DNA Banking for Plant Breeding, Biotechnology and Biodiversity Evaluation,” *Journal of Plant Research*, Vol.120, Issue.1, pp.17-29, 2007.
15. A. Karp, K.J. Edwards, M. Bruford, S. Funk, B. Vosman, M. Morgante, et al., “Molecular Technologies for Biodiversity Evaluation: Opportunities and Challenges,” *Nature Biotechnology*, Vol.15, Issue.7, pp.625-628, 1997.
16. K.C. Beng, R.T. Corlett, “Applications of Environmental DNA (eDNA) in Ecology and Conservation: Opportunities, Challenges and Prospects,” *Biodiversity and Conservation*, Vol.29, Issue.7, pp.2089-2121, 2020.
17. C.C. Azubuike, C.B. Chikere, G.C. Okpokwasili, “Bioremediation Techniques – Classification Based on Site of Application: Principles, Advantages, Limitations and Prospects,” *World Journal of Microbiology and Biotechnology*, Vol.32, Issue.11, Article 180, 2016.
18. H. Ali, E. Khan, M.A. Sajad, “Phytoremediation of Heavy Metals – Concepts and Applications,” *Chemosphere*, Vol.91, Issue.7, pp.869-881, 2013.
19. Shourav Dutta, “Effectiveness of Synthetic Biology to Diminish Global Biological Diversity Loss: A Comprehensive Review,” *International Journal of Scientific Research in Biological Sciences*, Vol.10, Issue.6, pp.31-38, 2023.
20. J. Schipper, J.S. Chanson, F. Chiozza, N.A. Cox, M. Hoffmann, V. Katariya, et al., “The Status of the World’s Land and Marine Mammals: Diversity, Threat, and Knowledge,” *Science*, Vol.322, Issue.5899, pp.225-230, 2008.
21. F. Forest, J. Moat, E. Baloch, N.A. Brummitt, S.P. Bachman, S. Ickert-Bond, et al., “Gymnosperms on the EDGE,” *Scientific Reports*, Vol.8, Article 6053, 2018.
22. A.C. Lees, L. Haskell, T. Allinson, S.B. Bezeng, I.J. Burfield, L.M. Renjifo, et al., “State of the World’s Birds,” *Annual Review of Environment and Resources*, Vol.47, pp.231-260, 2022.
23. N. Cox, B.E. Young, P. Bowles, M. Fernandez, J. Marin, G. Rapacciuolo, et al., “A Global Reptile Assessment Highlights Shared Conservation Needs of Tetrapods,” *Nature*, Vol.605, Issue.7909, pp.285-290, 2022.

24. K.E. Carpenter, M. Abrar, G. Aeby, R.B. Aronson, S. Banks, A. Bruckner, et al., “One-Third of Reef-Building Corals Face Elevated Extinction Risk from Climate Change and Local Impacts,” *Science*, Vol.321, Issue.5888, pp.560-563, 2008.
25. J. Clulow, S. Clulow, “Cryopreservation and Other Assisted Reproductive Technologies for the Conservation of Threatened Amphibians and Reptiles: Bringing the ARTs up to Speed,” *Reproduction, Fertility and Development*, Vol.28, Issue.8, pp.1116-1132, 2016.
26. M. Hagedorn, C.A. Page, K.L. O’Neil, D.M. Flores, L. Tichy, T. Conn, et al., “Assisted Gene Flow Using Cryopreserved Sperm in Critically Endangered Coral,” *Proceedings of the National Academy of Sciences of the USA*, Vol.118, Issue.38, Article e2110559118, 2021.
27. P. Buerger, C. Alvarez-Roa, C.W. Coppin, S.L. Pearce, L.J. Chakravarti, J.G. Oakeshott, O.R. Edwards, M.J.H. van Oppen, “Heat-Evolved Microalgal Symbionts Increase Coral Bleaching Tolerance,” *Science Advances*, Vol.6, Issue.20, Article eaba2498, 2020.